When all holes have the same length

Jake Horsfield^{*}, Myriam Preissmann[†], Cléophée Robin^{†‡}, Ni Luh Dewi Sintiari[‡], Nicolas Trotignon[‡] and Kristina Vušković^{*}

March 23, 2022

Abstract

For every integer $\ell \geq 7$, we give a structural description of the class of graphs whose chordless cycles of length at least 4 all have length ℓ .

1 Introduction

A hole in a graph is an induced cycle of length at least 4. For an integer $k \geq 7$, we study the class C_k of graphs where every hole has length k. Note that when k is even, this is a class of perfect graphs, and when k is odd, this is a class of even-hole-free graphs. Both these classes are well studied and we do not recall their definition. They have celebrated decomposition theorems (see [4] and [9]), but no full structural description. This motivates studying C_k .

In [16], the class of $(4K_1, C_4, C_6, C_7)$ -free graphs is studied. It is a subclass of C_5 . In [12], the class of $(4K_1, C_4, C_6)$ -free graphs is studied. In this class, every hole has length 5 or 7. In [2], the class of rings of length k

^{*}School of Computing, University of Leeds, UK and Faculty of Computer Science (RAF), Union University, Belgrade, Serbia. Partially supported by EPSRC grant EP/N0196660/1, and Serbian Ministry of Education and Science projects 174033 and III44006. sc15jh@leeds.ac.uk (Jake Horsfield), k.vuskovic@leeds.ac.uk

[†]Univ. Grenoble Alpes, CNRS, Grenoble INP, G-SCOP, 38000 Grenoble, France. myriam.preissmann@grenoble-inp.fr, cleophee.robin@grenoble-inp.fr

[‡]Univ Lyon, EnsL, UCBL, CNRS, LIP, F-69342, LYON Cedex 07, France. Partially supported by the LABEX MILYON (ANR-10-LABX-0070) of Université de Lyon, within the program "Investissements d'Avenir" (ANR-11-IDEX-0007) operated by the French National Research Agency (ANR) and by Agence Nationale de la Recherche (France) under research grant ANR DIGRAPHS ANR-19-CE48-0013-01. nld.sintiari@gmail.com, nicolas.trotignon@ens-lyon.fr

is defined for every integer $k \ge 4$ (see Section 3.2 for the definition), and it is used as a basic class for several decompositions theorems. Rings of length k form a subclass of C_k . In [15], a polynomial time algorithm that colors every ring is given. In [14], it is proved that for every fixed integer k, there exist rings of length k of arbitrarily large rankwidth.

In [1] and [3], polynomial-time algorithms that, given a graph G and $u, v \in V(G)$, decide whether there exists a path from u to v that is not a shortest path are described. It is easy to deduce from such algorithms an algorithm to recognize C_k in polynomial time.

Here, we provide first a structural description of graphs in $C_{2\ell+1}$ for any $\ell \geq 3$. It says that every graph in the class is constructed in some precise way or has a universal vertex or has a clique cut. The formal statement is given in Theorem 7.1. This work appears in the PhD thesis of Cléophée Robin [17]. Part of it and algorithmic applications will appear in the PhD thesis of Jake Horsfield. In the second part of this work, we provide a similar description of graphs in $C_{2\ell}$ for any $\ell \geq 4$. The formal statement is given in Theorem 10.1.

A similar description was obtained independently by Linda Cook and Paul Seymour. Much of it forms part of the PhD thesis [7] of Linda Cook and both groups decided to write a joint work based on this version, see [8]. Our statement is different but equivalent to the one in [8] as will be shown in the PhD thesis of Jake Horsfield. We publish the present version as a preprint because the approach is not the same and for later reference.

2 Definition and notation

We denote by \overline{G} the complement of a graph G.

When x is a vertex of a graph G and A is a subset of vertices of G or an induced subgraph of G, we denote by $N_A(x)$ the set of neighbors of x that are in A. Note that $x \notin N_A(x)$. We set $N_A[x] = \{x\} \cup N_A(x)$. If $X \subseteq V(G)$, we set $N_A(X) = (\bigcup_{x \in X} N_A(x)) \setminus X$ and $N_A[X] = N_A(X) \cup X$. We sometimes write N instead of $N_{V(G)}$ (when there is no risk of confusion).

A set $X \subseteq V(G)$ is complete to a set $Y \subseteq V(G)$ if they are disjoint and every vertex of X is adjacent to every vertex of Y. A set $X \subseteq V(G)$ is anticomplete to a set $Y \subseteq V(G)$ if they are disjoint and no vertex of X is adjacent to a vertex of Y. We sometimes say that x is complete (resp. anticomplete) to Y to mean that $\{x\}$ is complete (resp. anticomplete) to Y.

A vertex v in a graph G is *isolated* if it has no neighbors in G. It is *universal* if it is adjacent to all vertices of $G \setminus v$. A graph G is *connected* if

for every pair of vertices u, v there exists a path from u to v in G. A graph is *anticonnected* if its complement is connected. A *connected component* of a graph G is a subset X of V(G such that G[X] is connected and X is maximal w.r.t. this property. An *anticonnected component* of a graph G is a subset X of V(G) such that G[X] is anticonnected and X is maximal w.r.t. this property.

We will use the notion of *hypergraph*; that is, a structure similar to graphs except that the edges (called *hyperedges*) may contain an arbitrary positive number of vertices. While all the graphs that we use are simple, in hypergraphs, we allow hyperedges that contain a single vertex and multiple hyperedges (that is, there can be different hyperedges on the same set of vertices). Observe that we do not allow an empty hyperedge.

A cutset in a graph G is a set S of vertices such that $G \setminus S$ is disconnected. A clique in a graph is a set of pairwise adjacent vertices. In a graph, we view the empty set as a clique, and as a clique cutset of any disconnected graph. A stable set in a graph is a set of pairwise non-adjacent vertices.

For $k \geq 1$, we denote by P_k the path on k vertices, that is, the graph with vertex-set $\{p_1, \ldots, p_k\}$ and edge-set $\{p_1p_2, \ldots, p_{k-1}p_k\}$. We denote it by $p_1p_2 \ldots p_k$. If $1 \leq i \leq j \leq k$, we then denote by p_iPp_j the path $p_ip_{i+1} \ldots p_j$. For $k \geq 3$, we denote by C_k the cycle on k vertices; that is, the graph with vertex-set $\{p_1, \ldots, p_k\}$ and edge-set $\{p_1p_2, \ldots, p_{k-1}p_k, p_kp_1\}$. We denote it by $p_1p_2 \ldots p_kp_1$. We denote it by $p_1p_2 \ldots p_kp_1$. When C_k is a subgraph of a graph G (possibly not induced), an edge with both ends in $\{p_1, \ldots, p_k\}$ that is not an edge of C_k is called a *chord* of C_k . We denote by $2K_2$ the complement of C_4 .

We say that P is a path in a graph G (or P is a path of G) to mean that P is a path that is an induced subgraph of G. A hole in a graph G is a cycle of length at least 4 that is an induced subgraph of G. The length of a path, cycle or hole is the number of its edges. A hole is even or odd depending on the parity of its length.

A graph G contains a graph H if H is isomorphic to an induced subgraph of G and G is H-free if G does not contain H. For a class of graphs \mathcal{H} , we say that G is \mathcal{H} -free, if G is H-free for all H in \mathcal{H} .

3 A survey of some classes of graphs

Here we present several known classes of graphs and their properties. We do not need all of them, but we believe that presenting them all gives a better understanding of the class we work on.

3.1 Classes of perfect graphs

A graph is chordal if it is hole-free. A graph is a cograph if it is P_4 -free. A graph is a split graph if it is $(C_4, C_5, 2K_2)$ -free. A graph is a quasithreshold graph if it is (P_4, C_4) -free (quasi-threshold graphs are sometimes called trivially perfect graphs, see [13]). A graph is a threshold graph if it is $(P_4, C_4, 2K_2)$ -free (threshold graphs are sometimes called graphs with Dilworth number 1). A graph is a half graph if it is $(3K_1, C_4, C_5)$ -free.

Observe that these six classes are all classes of perfect graphs. The classes of cographs, split graphs and threshold graphs are self-complementary while the classes of chordal graphs, quasi-threshold and half graphs are not. In Figure 1, a Venn diagram of seven graph classes is represented (chordal and quasi – threshold mean complements of chordal and quasi-threshold graphs respectively). In every set, a typical example of the class is represented. The diagram provides several alternative definitions of the classes we work on (for instance, a threshold graph is a split cograph, a split graph is a chordal graph whose complement is chordal, and so on). All the information given by Figure 1 is easily recovered from the definitions of the corresponding classes.

Theorem 3.1 ([10]) A graph G is chordal if and only if every noncomplete induced subgraph of G has a clique cutset.

Theorem 3.2 ([18]) A graph G is a cograph if and only if every induced subgraph of G on at least two vertices is either not connected or not anticonnected.

Theorem 3.3 ([11]) A graph G is a split graph if and only if V(G) can be partitioned into a (possibly empty) clique and a (possibly empty) stable set.

The line graph of a hypergraph \mathcal{H} is the graph G whose vertex-set is $E(\mathcal{H})$ and where two hyperedges of \mathcal{H} are adjacent vertices of G whenever their intersection is non-empty. Recall that in this paper, hypergraphs may have multiple hyperedges (that are distinct hyperedges with the same vertices in them). A hypergraph is *laminar* is for every pair X, Y of hyperedges, either $X \subseteq Y$ or $Y \subseteq X$ or $X \cap Y = \emptyset$.

Theorem 3.4 ([20]) For all graphs G the following statements are equivalent.

(a) G is a quasi-threshold graph.



Figure 1: Venn diagram of seven classes of graphs

- (b) Every induced subgraph of G is disconnected or has a universal vertex.
- (c) G is the line graph of a laminar hypergraph.

Definition 3.5 If u and v are vertices of a graph G we write $u \leq_G v$ if $N(u) \setminus \{v\} \subseteq N(v) \setminus \{u\}$ and $u <_G v$ if $N(u) \setminus \{v\} \subsetneq N(v) \setminus \{u\}$.

Lemma 3.6 The relations \leq_G and \leq_G are transitive, that is, for all vertices u, v, w of some graph G, if $u \leq_G v$ and $v \leq_G w$, then $u \leq_G w$ (resp. if $u <_G v$ and $v <_G w$, then $u <_G w$).

We define \geq_G and $>_G$ accordingly (i.e. $x \geq_G y$ if and only if $y \leq_G x$) and extend these relations to sets of vertices X and Y as follows: $X \leq_G Y$ if and only if for every $x \in X$ and $y \in Y$, $x \leq_G y$ and so on.

Theorem 3.7 ([5]) For all graphs G the following statements are equivalent.

- (a) G is a threshold graph.
- (b) Every induced subgraph of G has an isolated vertex or a universal vertex.
- (c) For all vertices u and v of G, $u \leq_G v$ or $v \leq_G u$.

It is convenient to sort the vertices of a threshold graph. Formally, an ordering v_1, \ldots, v_k such that $v_i \leq_G v_j$ for all integers i and j satisfying $1 \leq i \leq j \leq k$ is called a *domination ordering*. There is another convenient ordering of the vertices of a threshold graph. By characterization (b) in Theorem 3.7, every threshold graph can be obtained by the following inductive process: start with a vertex u_1 , assume for some $k \geq 1$ that vertices u_1, \ldots, u_k are already constructed, and then add a vertex u_{k+1} that is either complete or anticomplete to $\{u_1, \ldots, u_k\}$. The order u_1, \ldots, u_n is then called an *elimination ordering* of the threshold graph (and it is not a domination ordering in general).

An example is represented in Figure 2. On the top, a threshold graph J on $\{v_1, \ldots, v_{10}\}$ is represented for which (v_1, \ldots, v_{10}) is a domination ordering. Vertices are circles with a number in them that gives the place of the vertex in the elimination ordering. On the bottom, the complement J' of J is represented. It is also a threshold graph but the domination ordering is reversed (it is (v_{10}, \ldots, v_1)), while the elimination ordering remains the same.



Figure 2: A threshold graph and its complement

Theorem 3.8 (Folklore) A graph G is a half graph if and only if V(G) can be partitioned into two (possibly empty) cliques K and K' such that for all vertices x and y in K (resp. in K'), either $x \leq_G y$ or $y \leq_G x$.

Proof. If G is a half graph, then the complement of G contains (as a subgraph, not necessarily induced) no cycle of odd length because a shortest such cycle cannot have length 3 (it would yield a $3K_1$ in G), cannot have length 5 (it would yield a C_5 in G) and cannot have length at least 7 (it would yield a C_4 in G). It follows that the complement of G is a bipartite graph, so V(G) can be partitioned into two cliques as claimed. The condition on \leq_G then follows from the fact that G contains no C_4 .

The converse statement is clear.

3.2 Classes defined by excluding Truemper configurations

Truemper configurations are graphs that play a role in many decomposition theorems, see [19]. They are the prisms, thetas, pyramids and wheels. Let us define them.

A prism is a graph made of three vertex-disjoint paths $P_1 = a_1 \dots b_1$, $P_2 = a_2 \dots b_2$, $P_3 = a_3 \dots b_3$ of length at least 1, such that $a_1 a_2 a_3$ and $b_1 b_2 b_3$ are triangles and no edges exist between the paths except those of the two triangles.

A pyramid is a graph made of three paths $P_1 = a \dots b_1$, $P_2 = a \dots b_2$, $P_3 = a \dots b_3$ of length at least 1, two of which have length at least 2, vertexdisjoint except at a, and such that $b_1b_2b_3$ is a triangle and no edges exist between the paths except those of the triangle and the three edges incident to a. The vertex a is called the *apex* of the pyramid.

A theta is a graph made of three internally vertex-disjoint paths $P_1 = a \dots b$, $P_2 = a \dots b$, $P_3 = a \dots b$ of length at least 2 and such that no edges exist between the paths except the three edges incident to a and the three edges incident to b.

Observe that the lengths of the paths used in the three definitions above are designed so that the union of any two of the paths induce a hole. A prism, pyramid or theta is *balanced* if the three paths in the definition are of the same length. It is *unbalanced* otherwise.



Figure 3: Pyramid, prism, theta and wheel (dashed lines represent paths)

A wheel W = (H, c) is a graph formed by a hole H (called the *rim*) together with a vertex c (called the *center*) that has at least three neighbors in the hole.

A wheel is a *universal wheel* if the center is adjacent to all vertices of the rim. A wheel is a *twin wheel* if the center is adjacent to exactly three vertices of the rim and they induce a P_3 . A wheel is *proper* if it is neither a twin wheel nor a universal wheel.

Truemper configurations are of interest here because of the following easy observation.

Lemma 3.9 Every unbalanced prism, every unbalanced pyramid, every unbalanced theta and every proper wheel contains holes of different lengths.

Every pyramid contains an odd hole. Every prism and every theta contains an even hole.

Proof. In a prism, pyramid or theta, the union of any two paths used in the definition induces a hole. Paths of different lengths are then easily used to provide holes of different lengths. In a proper wheel, the rim and a shortest hole are holes of different lengths.

In a pyramid, paths of the same parity, that exist since there are three paths, induce an odd hole. In thetas and prisms, they induce an even hole. \Box

The following variant is more useful for our study.

Lemma 3.10 If $\ell \geq 2$ is an integer and $G \in C_{2\ell+1}$, then every Truemper configuration of G is a twin wheel, a universal wheel or a pyramid whose three paths all have length ℓ .

If $\ell \geq 2$ is an integer and $G \in C_{2\ell}$, every Truemper configuration of G is a twin wheel, a universal wheel, a theta whose three paths all have length ℓ or a prism whose three paths all have length $\ell - 1$.

Proof. Clear from Lemma 3.9.

A graph G is *universally signable* if G is (prism, pyramid, theta, wheel)-free.

Theorem 3.11 ([6]) A graph G is universally signable if and only if every induced subgraph of G is a hole, a complete graph or has a clique cutset.

A graph G is a ring if its vertex-set can be partitioned into $k \ge 4$ sets K_1, \ldots, K_k such that (with subscripts understood to be taken modulo k):

- (a) K_1, \ldots, K_k are cliques;
- (b) for all $i \in \{1, \ldots, k\}$, K_i is anticomplete to $V(G) \setminus (K_{i-1} \cup K_i \cup K_{i+1})$;
- (c) for all $i \in \{1, \ldots, k\}$, some vertex of K_i is complete to $K_{i-1} \cup K_{i+1}$;
- (d) for all $i \in \{1, ..., k\}$ and all $x, x' \in K_i$, either $x \leq_G x'$ or $x' \leq_G x$.

The integer k in the definition above is the *length of the ring*. Observe that when $k \ge 4$, the hole C_k is a ring of length k. Observe also that, by Theorem 3.8, for any integer $1 \le i \le k$, the graph $G[K_i \cup K_{i+1}]$ is a half graph. We refer to the cliques K_1, \ldots, K_k as the *cliques of the ring* G.

Lemma 3.12 Every hole in a ring G of length k has length k.

Proof. We prove that a hole C in G contains at most one vertex in each clique of the ring. Suppose otherwise. Let $x, x' \in K_i$ be two vertices of C and suppose up to symmetry that $x \leq_G x'$. Hence, the neighbor of x in $C \setminus x'$ is also adjacent to x', so C contains a triangle, a contradiction.

Hence, C contains exactly one vertex in each clique of the ring. So, it has length k.

The following is a corollary of Theorem 1.6 from [2].

Theorem 3.13 If G is (prism, theta, pyramid, proper wheel, C_4 , C_5)-free, then one of the following holds.

- (a) G is a ring of length at least 6;
- (b) G has a clique cutset;
- (c) G has a universal vertex.

4 Odd templates

Here we define and study the main basic class of Theorem 7.1.

4.1 Modules in threshold graphs

Let G be a graph. A module of G is a set $X \subseteq V(G)$ such that every vertex in $V(G) \setminus X$ is either complete or anticomplete to X. Observe that all subsets of V(G) of cardinality 0, 1 or |V(G)| are modules of G. We will use the notion of module only in the context of threshold graphs. The reader can check that sets of vertices that are intervals for both elimination and domination orderings are modules. We omit the proof since we do not need this formally. We now state three lemmas.

Lemma 4.1 Let J be a threshold graph and $X \subseteq V(J)$ such that $|X| \ge 2$. Then \overline{J} is a threshold graph, X is a module of J if and only if it is a module of \overline{J} , and exactly one of J[X] and $\overline{J}[X]$ is anticonnected.

Proof. Being a threshold graph and module are properties that are closed under taking the complement. By Theorem 3.7, exactly one of J[X] or $\overline{J}[X]$ contains an isolated vertex, and the other one contains a universal vertex. Hence, since $|X| \ge 2$, exactly one of J[X] or $\overline{J}[X]$ is connected and the other one is anticonnected.

Lemma 4.2 Let J be a threshold graph. If X is an anticonnected module of J that contains at least two vertices, then N(X) is a clique that is complete to X. Moreover, $N(X) >_J X$.

Proof. Since X is a module, N(X) is complete to X. Suppose that N(X) is not a clique and let u and v be two non-adjacent vertices in N(X). Since $|X| \ge 2$ and X is anticonnected, X contains two non-adjacent vertices u', v'

that together with u and v form a C_4 in J. This contradicts J being a threshold graph.

Suppose that $N(X) >_J X$ does not hold. So, there exists $u \in N(X)$ and $v \in X$ with $v \ge_J u$. Since $u \in N(X)$, u is complete X, so v is complete to $X \setminus \{v\}$. This contradicts X being anticonnected.

Lemma 4.3 Let J be a threshold graph and $X \subseteq V(J)$ a module of J. If X contains some isolated vertices of J, then either X contains only isolated vertices of J, or X contains all non-isolated vertices of J.

Proof. Let S be the set of all isolated vertices of J and $T = V(J) \setminus S$. By assumption, X contains a vertex of S. If X contains only vertices of S, then the conclusion holds, so suppose that X contains at least one vertex of T. Suppose for a contradiction that X does not contain all of T. Since J[T] is connected (because it is a threshold graph with no isolated vertices and hence by Theorem 3.7 it contains a universal vertex), there exists an edge uv of J with $u \in T \cap X$ and $v \in T \setminus X$. Since X contains isolated vertices, this contradicts X being a module.

4.2 Templates

For an integer $\ell \geq 2$, an *odd* ℓ -*template* is any graph G that can be built according to the following process.

- (a) Choose a threshold graph J on vertex set $\{1, \ldots, k\}, k \ge 3$.
- (b) Choose a laminar hypergraph \mathcal{H} on vertex set $\{1, \ldots, k\}$ such that:
 - 1. every hyperedge X of \mathcal{H} is a module of J of cardinality at least 2 and
 - 2. at least one hyperedge W of \mathcal{H} contains all vertices of \mathcal{H} .
- (c) For each $i \in \{1, \ldots, k\}$, G contains two vertices v_i and v'_i that are linked by a path of G of length $\ell 1$. The k paths built at this step are vertex disjoint and are called the *principal paths* of the odd template.
- (d) The set of vertices of G is $V(G) = A \cup A' \cup B \cup B' \cup I$ where:
 - 1. I is the set of all internal vertices of the principal paths,
 - 2. $A = \{v_1, \ldots, v_k\},\$

- 3. $A' = \{v'_1, \dots, v'_k\},\$
- 4. $B = \{v_X : X \text{ hyperedge of } \mathcal{H} \text{ such that } J[X] \text{ is anticonnected} \},$
- 5. $B' = \{v'_X : X \text{ hyperedge of } \mathcal{H} \text{ such that } \overline{J}[X] \text{ is anticonnected} \}.$

Note that by Lemma 4.1, for every hyperedge X of \mathcal{H} , either $v_X \in B$ or $v'_X \in B'$ (and not both).

- (e) The set of edges of G is defined as follows.
 - 1. for every $v_i, v_j \in A$, $v_i v_j \in E(G)$ if and only if $ij \in E(J)$,
 - 2. for every $v'_i, v'_j \in A', v'_i v'_j \in E(G)$ if and only if $ij \notin E(J)$,
 - 3. for every $v_X, v_Y \in B, v_X v_Y \in E(G)$ if and only if $X \cap Y \neq \emptyset$,
 - 4. for every $v'_X, v'_Y \in B', v'_X v'_Y \in E(G)$ if and only if $X \cap Y \neq \emptyset$,
 - 5. for every $v_i \in A$, $v_X \in B$, $v_i v_X \in E(G)$ if and only if $i \in N_J[X]$,
 - 6. for every $v'_i \in A', v'_X \in B', v'_i v'_X \in E(G)$ if and only if $i \in N_{\bar{J}}[X]$,
 - 7. for every $v \in I$, v is incident to exactly two edges (those in its principal path).

The following notation is convenient.

Notation: For every vertex $x \in B$ such that $x = v_X$ where X is a hyperedge of \mathcal{H} , we set $H_x = \{v_i : i \in X\}$. Similarly, for every vertex $x \in B'$ such that $x = v'_X$ where X is a hyperedge of \mathcal{H} , we set $H'_x = \{v'_i : i \in X\}$.

We now list some properties of templates that follow directly from the definition.

- (i) G[A] is a threshold graph isomorphic to J and G[A'] is a threshold graph isomorphic to \overline{J} (and hence to the complement of G[A]).
- (ii) For all $x \in B$, H_x is a module of G[A] and $G[H_x]$ is anticonnected. Also for all $x \in B'$, H'_x is a module of G[A'] and $G[H'_x]$ is anticonnected.
- (iii) G[B] is isomorphic to the line graph of the hypergraph \mathcal{H}_B on vertex set A and hyperedge set $\{H_x : x \in B\}$. Also G[B'] is isomorphic to the line graph of the hypergraph $\mathcal{H}_{B'}$ on vertex set A' and hyperedge set $\{H'_x : x \in B'\}$. Hence G[B] and G[B'] are a quasi-threshold graphs by Theorem 3.4.
- (iv) There is an edge between $v_i \in A$ and $x \in B$ if and only if $v_i \in N_A[H_x]$, and there is an edge between $v'_i \in A'$ and $x \in B'$ if and only if $v'_i \in N_{A'}[H'_x]$.

Lemma 4.4 There exist vertices w and w' that are universal vertices in respectively $G[A \cup B]$ and $G[A' \cup B']$, and such that either $w \in A$ and $w' \in B'$, or $w \in B$ and $w' \in A'$.

Proof. By Theorem 3.7, G[A] contains a vertex u that is either universal or isolated. If u is universal, then for every $x \in B$, $u \in N(H_x)$ (u cannot be in H_x since $G[H_x]$ is anticonnected by property (ii) of templates). So, u is adjacent to x by property (iv) of templates. Hence, u is a universal vertex of $G[A \cup B]$.

Otherwise, u is an isolated vertex of G[A]. So, G[A] is anticonnected. Hence, the vertex w corresponding to the hyperedge W from condition (b2) of templates is in B. By property (iv) of templates, w is a universal vertex of $G[A \cup B]$.

The proof for $G[A' \cup B']$ is similar. So, w and w' exist, and by the way we construct them, we see that either $w \in A$ and $w' \in B'$, or $w \in B$ and $w' \in A'$.

Let w and w' be as in Lemma 4.4. The 7-tuple (A, B, A', B', I, w, w') is then called an ℓ -partition of G.

Let us give a simple example. Consider an integer $\ell \geq 2$ and a threshold graph J on three vertices $\{1, 2, 3\}$ with no edges. So, G[A] has no edges, G[A'] is a triangle on three vertices v'_1, v'_2, v'_3 , and for i = 1, 2, 3, there is a path of length $\ell - 1$ from v_i to v'_i . Consider \mathcal{H} the hypergraph on $\{1, 2, 3\}$ with a unique hyperedge that is $\{1, 2, 3\}$. We now see that G is a balanced pyramid with apex w and triangle $v'_1v'_2v'_3$. Under these circumstances, the sets $A = \{v_1, v_2, v_3\}, B = \{w\}, A' = \{v'_1, v'_2, v'_3\}, B' = \emptyset, I = V(G) \setminus (A \cup B \cup A' \cup B'), w$ and v'_3 form an ℓ -partition of G.

It is worth noting that the ℓ -partition above is not unique. Here is another one. Call u the neighbor of v'_3 in the path from v_3 to v'_3 with interior in I (possibly, $u = v_3$). Set $A_1 = \{w, v_1, v_2\}$, $B_1 = \emptyset$, $A'_1 = \{u, v'_1, v'_2\}$, $B'_1 = \{v'_3\}$ and $I_1 = V(G) \setminus (A_1 \cup B_1 \cup A'_1 \cup B'_1)$. It can be checked that A_1 , B_1, A'_1, B'_1, I_1, w and v'_3 form another ℓ -partition of G. See Figure 4. Some edges are dashed in several ways, this will be explained later, so far, they are just edges of G.

Lemma 4.5 For all integers $\ell \geq 2$, every pyramid Π such that $\Pi \in C_{2\ell+1}$ is an odd ℓ -template.

Proof. Since $\Pi \in C_{2\ell+1}$, its three paths have length ℓ . The explanations above show it is an odd ℓ -template.



Figure 4: Two 3-partitions of a pyramid whose paths all have length 3

We now give a more complicated example represented in Figure 5. The threshold graph J has 10 vertices. Each vertex of G[A] and G[A'] is represented with a number in a circle that represents the elimination ordering of the threshold graph it belongs to. The hypergraph \mathcal{H} has the following hyperedges: $X_1 = \{1, 2\}, X_2 = \{1, 2, 3\}, X_3 = \{9, 10\}, X_4 = \{5, 6, 7\}, X_5 = \{5, 6, 7, 8\}, X_6 = \{4, 5, 6, 7, 8\}$ and $X_7 = \{1, \ldots, 10\}$. The vertex of $B \cup B'$ corresponding to a hyperedge X_i is denoted by x_i .

4.3 Structure of odd templates

Throughout this subsection, $\ell \geq 2$ is an integer and (A, B, A', B', I, w, w') is an ℓ -partition of an odd ℓ -template G.

Lemma 4.6 If $x \in B$ (resp. $x \in B'$), then H_x (resp. H'_x) is the unique anticomponent of $G[N_A(x)]$ (resp. $G[N_{A'}(x)]$) that contains at least two vertices.

Proof. Since G is C_4 -free and $N_A(x)$ contains at least two non-adjacent vertices, $G[N_A(x)]$ contains a unique anticomponent X of size at least 2. Since by property (ii) of templates, H_x is an anticonnected module of G[A], it is also an anticonnected module of $G[N_A(x)]$. Since every vertex of $N_A(x)$ is either in H_x or complete to H_x , H_x must be an anticomponent of $G[N_A(x)]$, and since it contains at least two vertices, it is equal to X. \Box



Figure 5: An odd 4-template

Lemma 4.7 If $x, y \in B$ (resp. $\in B'$) are such that $xy \notin E(G)$, then $H_x \cup \{x\}$ (resp. $H'_x \cup \{x\}$) is anticomplete to $H_y \cup \{y\}$ (resp. $H'_y \cup \{y\}$).

Proof. Suppose $x, y \in B$ and $xy \notin E(G)$. Then by condition (e3) of templates, H_x and H_y are disjoint.

Suppose there is at least one edge from H_x to H_y . Since they are both modules of G[A], it follows that H_x is complete to H_y , so $H_y \subseteq N(H_x)$. Hence, by Lemma 4.2, H_y is a clique. Since H_y contains at least two vertices, this contradicts H_y being anticonnected. So, H_x is anticomplete to H_y . Hence, by property (iv) of templates, x is anticomplete to H_y and y is anticomplete to H_x . So, $H_x \cup \{x\}$ is anticomplete to $H_y \cup \{y\}$ because $xy \notin E(G)$ holds from our assumption.

The proof for $x, y \in B'$ is similar.

 \mathbf{L}

Lemma 4.8 Every vertex of G has degree at least 2 and every vertex of
$$B \cup B'$$
 has degree at least 3.

Proof. Vertices in I are all in the interior of some path, so they have degree at least 2.

Vertex w has degree at least 2 since $|A| \geq 3$. A vertex $v \in A \setminus \{w\}$ therefore has degree at least 2 (one neighbor in I, and w). So every vertex of A has degree at least 2. The proof for A' is similar.

Let x be a vertex of B. If x = w, then x has degree at least 3 (because $|A| \geq 3$), so we may assume $x \neq w$. By, property (ii) of templates, $|H_x| \geq 2$ and $w \notin H_x$ because H_x is anticonnected. So x has degree at least 3 as claimed (at least two neighbors in H_x , and w). The proof for $x \in B'$ is similar.

The following shows that odd templates can be considered as a generalization of balanced pyramids (we do not need it and include it because we believe it helps understanding the structure of the class we work on).

Lemma 4.9 For every integer $\ell \geq 2$, every odd ℓ -template G contains a pyramid.

Proof. Consider three vertices v_i, v_j and v_h in A and the corresponding vertices v'_i, v'_j and v'_h in A'. Exactly one of $G[\{v_i, v_j, v_h\}]$ and $G'[\{v'_i, v'_j, v'_h\}]$ is connected (because they have three vertices and one is isomorphic to the complement of the other). So, up to symmetry, we may assume that $G[\{v_i, v_j, v_h\}]$ is disconnected (and therefore contains at most one edge).

Note that w is distinct from v_i, v_j and v_h since $G[\{v_i, v_j, v_h\}]$ is disconnected. We see that w and the three principal paths linking $\{v_i, v_j, v_h\}$ to $\{v'_i, v'_j, v'_h\}$ form a pyramid (if $G[\{v_i, v_j, v_h\}]$ contains one edge e, then the triangle is formed by e and w, and otherwise it is $v'_i v'_j v'_h$).

From the definition of odd ℓ -templates, every vertex $x \in B$ corresponds to a set $H_x \subseteq A$. These sets form a hypergraph \mathcal{H}_B on the vertex-set A(that is isomophic to a sub-hypergraph of \mathcal{H}). Let us build an extention \mathcal{H}_A of \mathcal{H}_B by adding more hyperedges: for every vertex $v \in A$, we add the hyperedge

$$H_v = N_A[v] \cap \{ u \in A : u \leq_{G[A]} v \}.$$

Note that $v \in H_v$.

Lemma 4.10 \mathcal{H}_A is a laminar hypergraph and $G[A \cup B]$ is isomorphic to its line graph (in particular, $G[A \cup B]$ is a quasi-threshold graph and therefore a chordal graph). A similar statements holds for $G[A' \cup B']$.

Proof. By construction, every vertex of $A \cup B$ corresponds to a hyperedge of \mathcal{H}_A . We have to check that the ends of every edge of $G[A \cup B]$ correspond to hyperedges of \mathcal{H}_A that are included one in the other, and that the ends of every non-edge correspond to a pair of disjoint hyperedges. This will prove that $G[A \cup B]$ is isomorphic to the line graph of \mathcal{H}_A and that \mathcal{H}_A is laminar. Let us check all the cases.

For $x, y \in B$, since \mathcal{H}_B is laminar and G[B] is isomorphic to its line graph, we have nothing to prove.

Let $u, v \in A$. By Theorem 3.7, we may assume up to symmetry that $u \geq_{G[A]} v$. If $uv \in E(G)$, then clearly $H_v \subseteq H_u$. Suppose $uv \notin E(G)$, and let t be a vertex of A such that $t \leq_{G[A]} v$. So, $t \leq_{G[A]} u$. If $tv \in E(G)$, then $uv \in E(G)$, a contradiction. So, $tv \notin E(G)$ and $H_v = \{v\}$. Since $v \notin N[u]$, we have that $H_u \cap H_v = \emptyset$.

Consider finally vertices $u \in A$ and $x \in B$. Suppose first that $ux \in E(G)$. By property ((iv)) of templates, we have that $u \in N_A[H_x]$. If $u \in H_x$, then $H_u \subseteq H_x$ by Lemma 4.2 (specifically, we use $N_A(H_x) >_{G[A]} H_x$ to conclude that $N_A(H_x) \cap H_u = \emptyset$, and then since $H_u \subseteq N_A[H_x]$ it follows that $H_u \subseteq H_x$). If $u \in N_A(H_x)$, then by Lemma 4.2 (again also using that $N_A(H_x) >_{G[A]} H_x$), $H_x \subseteq H_u$. So, an edge indeed yields an inclusion of the corresponding hyperedges.

Suppose now that $ux \notin E(G)$. So, $u \notin N[H_x]$. Since u is not in H_x and has no neighbor in H_x , it follows that H_u is disjoint from H_x .

So \mathcal{H}_A is a laminar hypergraph and $G[A \cup B]$ is isomorphic to its line graph. It follows from Theorem 3.4 that $G[A \cup B]$ is a quasi-threshold graph and therefore a chordal graph.

Lemma 4.11 Every hole of G is formed by two principal paths of G and a single vertex of $A \cup B \cup A' \cup B'$ that does not belong to these principal paths (it therefore has length $2\ell + 1$).

Proof. By Lemma 4.10, a hole C of G cannot contain only vertices of $A \cup B$, and similarly, it cannot contain only vertices of $A' \cup B'$. So it must contain vertices of some principal path, and also of a second principal path. In fact, C must go through exactly two principal paths, since G[A] is isomorphic to the complement of G[A'], if three paths are involved, there would be a vertex of C with three neighbors in C, a contradiction.

Since G[A] is isomorphic to the complement of G[A'], up to a symmetry, for some nonadjacent vertices $u, v \in A$, the hole C is made of a path $P = u \dots v$ with interior in $I \cup A'$ (whose length is $2\ell - 1$) and a path $Q = u \dots v$ of $G[A \cup B]$. By Lemma 4.10, Q has length at most 2 (because a quasithreshold graph is P_4 -free), and since $uv \notin E(G)$, it has length 2. So, C has length $2\ell + 1$ as claimed. \Box

4.4 Connecting vertices of a template

Lemma 4.12 If $x \in B$ and $y \in B'$, then there exists in G two paths P and Q of length $\ell+1$ from x to y such that P (resp. Q) contains a principal path P_0 (resp. Q_0), and $P_0 \neq Q_0$.

Proof. We set $X = \{i \in \{1, ..., k\} : v_i \in H_x\}$ and $Y = \{i \in \{1, ..., k\} : v'_i \in H'_y\}$. So, X and Y are hyperedges of \mathcal{H} and since \mathcal{H} is laminar, either $X \subseteq Y, Y \subseteq X$ or $X \cap Y = \emptyset$.

If $X \subseteq Y$, then let i, j be distinct members of X (and therefore of Y). The paths $xv_iP_iv'_iy$ and $xv_jP_jv'_jy$ are the paths we are looking for. The proof is similar when $Y \subseteq X$.

If $X \cap Y = \emptyset$, then let i, j, q, r be distinct integers such that $i, j \in X$ and $q, r \in Y$. Since G[A] is isomorphic to the complement of G[A'], we may assume up to symmetry that $v_i v_q \in E(G)$. So, $v'_i v'_q \notin E(G)$. Since H'_y is a module of G[A'], $v'_i v'_r \notin E(G)$. It follows that $v_i v_r \in E(G)$. So, $v_r, v_q \in N_A(H_x)$. Hence, by property (iv) of templates, $xv_r, xv_q \in E(G)$. It follows that $xv_q P_q v'_q y$ and $xv_r P_r v'_r y$ are the two paths we are looking for. \Box **Lemma 4.13** If $x \in A \cup B$ and $y \in A' \cup B'$, then there exists in G a path P of length $\ell - 1$, ℓ or $\ell + 1$ from x to y that contains a principal path. More specifically:

- If $x \in A$ and $y \in A'$, then P has length $\ell 1$ or ℓ .
- If $x \in A$ and $y \in B'$, or if $x \in B$ and $y \in A'$, then P has length ℓ or $\ell + 1$.
- If $x \in B$ and $y \in B'$, then P has length $\ell + 1$.

Proof. Suppose first that $x \in A$, say $x = v_i$. If $y \in A'$, then set $y = v'_j$. If i = j, then P_i has length $\ell - 1$. If $i \neq j$, then one of $v_i v_j P_j v'_j$ or $v_i P_i v'_i v'_j$ is a path of length ℓ . If $y \in B'$, then one of $v_i P_i v'_i y$ or $v_i P_i v'_i w' y$ is the path we are looking for. The proof is similar when $y \in A'$.

We may therefore assume that $x \in B$ and $y \in B'$. So one of the two paths obtained in Lemma 4.12 can be chosen.

4.5 Odd pretemplates

Checking that a graph is an odd ℓ -template is tedious. We now introduce a simpler notion that is in some sense equivalent. For every integer $\ell \geq 3$, an *odd* ℓ -*pretemplate* is a graph G whose vertex-set can be partitioned into five sets A, B, A', B' and I with the following properties.

- (a) $N(B) \subseteq A$ and $N(A \cup B) \subseteq I$.
- (b) $N(B') \subseteq A'$ and $N(A' \cup B') \subseteq I$.
- (c) $|A| = |A'| = k \ge 3$, $A = \{v_1, \dots, v_k\}$ and $A' = \{v'_1, \dots, v'_k\}$.
- (d) For every $i \in \{1, ..., k\}$, there exists a unique path P_i from v_i to v'_i whose interior is in I.
- (e) Every vertex in I has degree 2 and lies on a path from v_i to v'_i for some $i \in \{1, \ldots, k\}$.
- (f) All paths P_1, \ldots, P_k have length $\ell 1$.
- (g) $G[A \cup B]$ and $G[A' \cup B']$ are both connected graphs.
- (h) Every vertex of B is in the interior of a path of $G[A \cup B]$ with both ends in A.

(i) Every vertex of B' is in the interior of a path of $G[A' \cup B']$ with both ends in A'.

We then say that (A, B, A', B', I) is an ℓ -pretemplate partition of G. Note that templates are defined for all integers $\ell \geq 2$, while pretemplates are defined only when $\ell \geq 3$. In fact we do not need odd 2-templates, we defined them for possible later use.

It is easy to check that when $\ell \geq 3$, the five first elements of every ℓ -partition of G is an ℓ -pretemplate partition. The condition on the connectivity of $G[A \cup B]$ and $G[A' \cup B']$ follows Lemma 4.4. The condition (h) follows from the fact for every $x \in B$, H_x contains two non-adjacent vertices, so a vertex $x \in B$ lies on a path of length 2 with ends in A and condition (i) holds similarly. Conversely, we prove the following lemma (it is important to note that $\ell \geq 3$).

Lemma 4.14 Let $\ell \geq 3$ be an integer. If $G \in C_{2\ell+1}$ is an odd ℓ -pretemplate, then G is an odd ℓ -template. Moreover, for every odd ℓ -pretemplate partition (A, B, A', B', I) of G, there exist w and w' in V(G) such that (A, B, A', B', I, w, w') is an ℓ -partition of G.

Proof. Let (A, B, A', B', I) be an ℓ -pretemplate partition of G. We first study the structure of G[A] and G[A'].

(1) For all distinct $i, j \in \{1, ..., k\}$, $v_i v_j \in E(G)$ if and only if $v'_i v'_j \notin E(G)$. In particular, G[A] is isomorphic to the complement of G[A'].

If $v_i v_j, v'_i v'_j \in E(G)$, then P_i and P_j form a hole of even length, a contradiction. If $v_i v_j, v'_i v'_j \notin E(G)$, then P_i, P_j , a path from v_i to v_j in $G[A \cup B]$ and a path from v'_i to v'_j in $G[A' \cup B']$ form a hole of length at least $2\ell + 2$, a contradiction. This proves (1).

(2) Every path of $G[A \cup B]$ with both ends in A is of length at most 2.

Let $P = v_i \dots v_j$ be a path of $G[A \cup B]$ with both ends in A. If P has length at least 3, then by (1), paths P, P_i and P_j form a hole of length at least $2\ell + 2$, a contradiction. This proves (2).

(3) G[A] is a threshold graph.

G[A] is obviously C_4 -free. Since the complement of C_4 is $2K_2$ and since G[A'] is also C_4 -free, it follows by (1) that G[A] is $2K_2$ -free. By (2), G[A] is P_4 -free. So G[A] is $(P_4, C_4, 2K_2)$ -free and is therefore a threshold graph. This proves (3).

We now study the structure of G[B] and its relation with G[A].

(4) For every vertex $x \in B$, $G[N_A(x)]$ has a unique anticonnected component of size at least 2.

By the definition of odd pretemplates, x is in the interior of a path $P = v_i \dots v_j$ of $G[A \cup B]$ with both ends in A. By (2), P has length 2, so x is adjacent to v_i and v_j . Hence $G[N_A(x)]$ has an anticonnected component of size at least 2. It is unique, for otherwise G[A] contains a C_4 . This proves (4).

For all $x \in B$, we define H_x to be the anticonnected component of $G[N_A(x)]$ of size at least 2 whose existence follows from (4).

(5) For every x in B, H_x is a module of G[A].

Otherwise, since H_x is anticonnected and is not a module, there exists $v_h \in A \setminus H_x$ and non-adjacent $v_i, v_j \in H_x$ such that $v_i v_h \in E(G)$ and $v_j v_h \notin E(G)$. Note that $xv_h \notin E(G)$ because otherwise, v_h would be in H_x . Hence, v_i, x , P_j and P_h form a hole of length $2\ell + 2$, a contradiction. This proves (5).

(6) If xy is an edge of G[B], then $H_x \subseteq H_y$ or $H_y \subseteq H_x$.

Up to symmetry, we may assume that $N_A(x) \subseteq N_A(y)$, for otherwise vertices $v_i \in N_A(x) \setminus N_A(y)$ and $v_j \in N_A(y) \setminus N_A(x)$ either form a C_4 with x and y or a hole of length $2\ell + 2$ with P_i and P_j .

By (4), $G[N_A(y)]$ has only one anticonnected component of size at least 2, namely H_y . Since H_x is anticonnected, has size at least 2 and is included in $N_A(y)$, it must be included in H_y . This proves (6).

(7) If x and y are non-adjacent vertices of B, then H_x and H_y are disjoint.

On the contrary, suppose that x and y are nonadjacent vertices of B but there exists a vertex $v \in H_x \cap H_y$. Since H_x is anticonnected and of size at least 2, there exists $v_i \in H_x$ non-adjacent to v. Note that $v_i y \notin E(G)$, for otherwise x, y, v_i and v form a C_4 . Similarly, there exists a vertex $v_j \in H_y$ that is non-adjacent to v and to x. If $v_i v_j \in E(G)$, then $\{x, y, v, v_i, v_j\}$ induces a C_5 , a contradiction. Otherwise, P_i, P_j, x, y and v form a hole of length $2\ell + 3$, a contradiction. This proves (7).

We are now ready to define the hypergraph \mathcal{H} . For every $x \in B$, we defined a set $H_x \subseteq A$. We may define similarly a set $H'_x \subseteq A'$ for every $x \in B'$. From (6) and (7), the sets H_x for $x \in B$ form a laminar hypergraph \mathcal{H}_B (with vertex set A). Symetrically, the sets H'_x for $x \in B'$ form a laminar hypergraph $\mathcal{H}_{B'}$ (with vertex set A'). Let \mathcal{H} be the hypergraph whose vertex

set is $\{1, \ldots, k\}$ and such that $H \subseteq \{1, \ldots, k\}$ is a hyperedge of \mathcal{H} if and only if $H = \{i : v_i \in H_x\}$ for some $x \in B$ or $H = \{i : v'_i \in H'_x\}$ for some $x \in B'$.

(8) The hypergraph \mathcal{H} is laminar.

If \mathcal{H} is not laminar, then there exist $X, Y \in E(\mathcal{H})$ such that $X \setminus Y, Y \setminus X$ and $X \cap Y$ are all non-empty. Since \mathcal{H}_B and $\mathcal{H}_{B'}$ are both laminar, there exists $x \in B$ such that $H_x = \{v_i : i \in X\}$ and $y \in B'$ such that $H'_y = \{v'_i : i \in Y\}$.

We set $H_y = \{v_i : i \in Y\}$. Note that $H_x \setminus H_y$, $H_y \setminus H_x$ and $H_x \cap H_y$ are all non-empty. Also, because of the properties of H'_y and by (1), $G[H_y]$ is connected (because $G[H'_y]$ is anticonnected) and H_y is a module of G[A].

Since $G[H_x]$ is anticonnected, there exist non-adjacent vertices $u \in H_x \setminus H_y$ and $v \in H_x \cap H_y$. Since $G[H_y]$ is connected, there exists a path from v to $t \in H_y \setminus H_x$ and we may assume that vt is an edge. Since H_y is a module of G[A], $ut \notin E(G)$. So, t is adjacent to v and non-adjacent to u. This contradicts H_x being a module of G[A]. This proves (8).

We may now finish the proof of Lemma 4.14. We show how G can be built by the process described in the definition of odd templates. We start by setting $V(J) = \{1, \ldots, k\}$, and by making i adjacent to j in J if and only if $v_i v_j \in E(G)$. By (3), J is a threshold graph as required. Clearly condition (d) of odd ℓ -templates holds, the paths linking A to A' are as in condition (c) of odd ℓ -templates and condition (e7) of odd ℓ -templates holds. By (1), conditions (e1) and (e2) of odd ℓ -templates hold. We then consider the hypergraph \mathcal{H} defined above. It is laminar by (8). By (5), condition (b1) of templates is satisfied.

By definition of H_x , for every x in B, $N_A(x) \subseteq N_A[H_x]$. Suppose that there exists $u \in N_A[H_x] \setminus N_A(x)$. Since by (5) H_x is a module, it follows from Lemma 4.2 that u is complete to H_x , so x and u together with two nonadjacent vertices from H_x induce a C_4 , a contradiction. Hence, $N_A(x) = N_A[H_x]$ and condition (e5) of odd templates is satisfied.

By (6) and (7), condition (e3) of templates is satisfied.

By symmetry and by (1), conditions (e4) and (e6) of templates are satisfied. Therefore condition (e) of templates is satisfied.

To conclude the proof, let us check condition (b2) of templates. By (1), (3) and Theorem 3.7, up to symmetry, we may assume that G[A] contains an isolated vertex v_i . Since $G[A \cup B]$ is connected and $|A| \ge 3$ by the definition of odd pretemplates, there exists a path P in $G[A \cup B]$ from v_i to a vertex $u \in A \setminus \{v_i\}$. By (2) and since v_i has no neighbor in A, we have that $P = uyv_i$ where $y \in B$. So, H_y contains v_i . We may therefore consider the hyperedge W of \mathcal{H} that contains i and that is inclusion wise maximal w.r.t. this property. If there exists $j \in \{1, \ldots, k\} \setminus W$, since $v_j v_i \notin E(G)$, we deduce as above that \mathcal{H} has a hyperedge Z that contains i and j. Because of $j, Z \subseteq W$ is impossible; because of $i, W \cap Z = \emptyset$ is impossible; and because of the maximality of $W, W \subsetneq Z$ is impossible. Hence, W and Zcontradict \mathcal{H} being laminar. This proves that $W = \{1, \ldots, k\}$, as claimed in condition (b2) of templates.

Hence, $G[A \cup B]$ has universal vertex w. Also, $G[A' \cup B']$ has a universal vertex w' (we may apply Lemma 4.4 since we now know that G is an odd ℓ -template). So, (A, B, A', B', I, w, w') is an ℓ -partition of G.

4.6 Twins and proper partitions

Two distinct vertices x and y in a graph are twins if N[x] = N[y] (in particular, x and y are adjacent). A graph is twinless if it contains no twins.

Lemma 4.15 Let (A, B, A', B', I, w, w') be an ℓ -partition of an odd ℓ template G. Two vertices x and y of G are twins if and only if $x, y \in B$ and $H_x = H_y$, or $x, y \in B'$ and $H'_x = H'_y$.

Proof. If $x, y \in B$ and $H_x = H_y$, or $x, y \in B'$ and $H'_x = H'_y$, then x and y are obviously twins.

We claim that for all $x \in A \cup I \cup A'$, there exist two vertices $a, b \in N_G(x)$ such that $N[a] \cap N[b] = \{x\}$. If $x \in I$ choose a and b to be the only two neighbors of x. If $x \in A$, then set a = w when $x \neq w$, and choose for a any vertex of $A \setminus \{x\}$ when x = w. Choose for b the neighbor of x in I. In both cases, by condition (e7) of templates, $N_G[a] \cap N_G[b] = \{x\}$. The proof is similar when $x \in A'$. So, x has no twin in G. \Box

An ℓ -partition (A, B, A', B', I, w, w') of an odd ℓ -template G is proper if one of G[A] or G[A'] contains at least two isolated vertices.

Lemma 4.16 For all integers $\ell \geq 3$, every twinless odd ℓ -template G admits a proper ℓ -partition.

Proof. Let (A, B, A', B', I, w, w') be an ℓ -partition of G such that the number M of isolated vertices of G[A] is maximum. We suppose that v_1, \ldots, v_k is a domination ordering of G[A].

By Theorem 3.7 and since we may swap A, B, w and A', B', w', by the maximality of M, v_1 is an isolated vertex of G[A]. It follows that $w \in B$.

By definition of templates, v'_1 is a universal vertex of $G[A' \cup B']$. Suppose for a contradiction that v_2 is not isolated in G[A]. So, M = 1.

Let H_x be any hyperedge of \mathcal{H}_B containing v_1 . By Lemma 4.3, since H_x contains a non-isolated vertex of G[A], it contains all of them. So, $H_x = A$. Hence, N[x] = N[w], so x = w since G is twinless. This proves that $N(v_1) = \{w, v_1^+\}$ where v_1^+ is the neighbor of v_1 in I. Let $v_1'^+$ be the neighbor of v_1' in I. We now describe a new partition of the vertices of G. We set:

- $A_1 = \{w, v_2, \dots, v_k\},\$
- $B_1 = B \setminus \{w\},$
- $A'_1 = \{v'^+_1, v'_2, \dots, v'_k\},\$
- $B'_1 = B' \cup \{v'_1\}$ and
- $I_1 = \{v_1\} \cup I \setminus \{v_1'^+\}.$

All conditions of the definition of a pretemplate are easily checked to be satisfied by $(A_1, B_1, A'_1, B'_1, I_1)$. By Lemma 4.11, every hole in Ghas length $2\ell + 1$. We may therefore apply Lemma 4.14 to prove that $(A_1, B_1, A'_1, B'_1, I_1, w, v'_1)$ is an ℓ -partition of G. So, $(A'_1, B'_1, A_1, B_1, I_1, v'_1, w)$ contradicts that maximality of M since $G[A'_1]$ has two isolated vertices, namely v'^+_1 and v'_k (note that since M = 1, it follows by Theorem 3.7 that $G[A \setminus \{v_1\}]$ has a universal vertex ; in particular v_k is a universal vertex of $G[A \setminus \{v_1\}]$ and hence v'_k is an isolated vertex of $G[A'_1]$). \Box

Lemma 4.17 Every proper ℓ -partition (A, B, A', B', I, w, w') of a twinless odd ℓ -template satisfies one of the following:

- $w \in B$, w is the unique universal vertex of $G[A \cup B]$, G[A] contains at least two isolated vertices, $H_w = A$, $w' \in A'$ and $A' \setminus \{w'\}$ contains at least one universal vertex of G[A'].
- w ∈ A, A \ {w} contains at least one universal vertex of G[A], w' ∈ B', w' is the unique universal vertex of G[A' ∪ B'], G[A'] contains at least two isolated vertices and H'_{w'} = A'.

Proof. Since (A, B, A', B', I, w, w') is a proper ℓ -partition, up to symmetry, we may assume that G[A] contains two isolated vertices. So, $w \in B$. By definition of ℓ -partitions, it follows that $w' \in A'$. Since G is twinless and

G[A] contains isolated vertices, w is the unique universal vertex of $G[A \cup B]$. By Lemma 4.6, $H_w = A$. Also $A' \setminus \{w'\}$ contains at least one universal vertex of G[A'] since G[A] contains two isolated vertices. \Box

We do not use the following lemma formally, but it illustrates a key property of proper partitions. In non-proper partitions, there may exist vertices in A that have degree 2 and have one neighbor in A and one in I. These are hard to think of, because they yield edges with both ends in A that can be "blown up" into a general half graph as we will see in the next section. The next lemma states that this situation does not occur with proper partitions.

Lemma 4.18 Suppose $\ell \geq 3$ and G is an odd ℓ -template with a proper ℓ -partition (A, B, A', B', I, w, w'). If a vertex v in $A \cup B \cup A' \cup B'$ has degree 2 (in G), then $v \in A \cup A'$ and v is adjacent to a vertex of $B \cup B'$ and has its other neighbor in I.

Proof. Up to symmetry, suppose that G[A] contains at least two isolated vertices. So, $w \in B$. Consider a vertex $v \in A \cup B \cup A' \cup B'$ that is of degree 2 in G. By Lemma 4.8, $v \notin B \cup B'$. Since G[A'] has two universal vertices, every vertex in A' has degree at least 3, so $v \in A$, and v is adjacent to $w \in B$ and to some vertex in I as claimed. \Box

5 Blowup

Our goal in this section is to see how a bigger graph can be obtained from a template G by turning every vertex into a non empty clique. This will be called *blowing up* G. In the blowup operation, non-adjacent vertices yield cliques that are anticomplete to each other. Adjacent vertices u and v yield cliques that are complete to each other in some situations (when uv is a so-called *solid edge* of the template), but in some other situations, they may yield pairs of cliques that induce a more general half graph, like when a ring is obtained from "blowing up" a chordless cycle. This happens when uvis a so-called *flat* or *optional edge* of the template. We now define all this formally.

Throughout all this section, $\ell \geq 3$ is an integer, G is a an odd ℓ -template with a fixed an ℓ -partition (A, B, A', B', I, w, w').

5.1 Flat, optional and solid edges

An edge of G is *flat* if at least one of its end is in I. An edge of G is *optional* if one end is a vertex $x \in B$ (resp. $x \in B'$) and the other end is a vertex $u \in H_x$ that is an isolated vertex of $G[H_x]$ (resp. a vertex $u \in H'_x$ that is an isolated vertex of $G[H_x]$). An edge that is neither flat nor optional is *solid*. See Figure 5 where solid edges are represented by solid lines, flat edges by doted lines and optional edges by dashed lines.

Observe that the status of an edge depends on the ℓ -partition of the odd ℓ -template. See Figure 4, where the same template is represented with two different ℓ -partitions. Recall that throughout this section, the ℓ -partition is fixed, and so is the status of the edges.

Lemma 5.1 If ux is an optional edge of G with $u \in A$ and $x \in B$, then $N_A(H_x) = N_A(u)$. Moreover, if $y \in B \setminus \{x\}$ and $yu \in E(G)$, then $H_x \subseteq H_y$ or $H_y \subseteq H_x$ (in particular, $xy \in E(G)$).

Proof. Since H_x is a module of G[A] and u is isolated in H_x , we have $N_A(H_x) = N_A(u)$. If the second conclusion fails, then since \mathcal{H}_B is laminar, $H_y \cap H_x = \emptyset$. So $xy \notin E(G)$. Since $yu \in E(G)$ and $u \notin H_y$, we have $u \in N_A(H_y)$, so u is complete to H_y since H_y is a module. So $H_y \subseteq N_A(u) = N_A(H_x)$, which is a clique by Lemma 4.2. This contradicts H_y being anticonnected.

A clique of G is *solid* if all its edges are solid.

Lemma 5.2 If ux is an optional edge of G such that $u \in A$ and $x \in B$, then $N_{A \cup B}(u)$ is a solid clique of G.

Proof. By Lemma 5.1, $N_A(H_x) = N_A(u)$. By Lemma 4.2, $N_A(H_x) = N_A(u)$ is a clique. It is solid because edges with both ends in A are solid. Hence $N_A(u)$ is a solid clique.

By Lemma 5.1 all vertices from $N_B(u)$ are adjacent since they correspond to hyperedges of \mathcal{H}_B that are included in each other. Therefore, $N_B(u)$ is a clique and it is solid because edges with both ends in B are solid. Hence $N_B(u)$ is a solid clique.

It remains to prove that $N_A(u)$ is complete to $N_B(u)$ and that all edges between these two sets are solid. So let $y \in B$ and $v \in A$ be two neighbors of u. Note that $v \notin H_x$ and possibly y = x. If $u \in H_y$, then vy is an edge because $v \in N_A(u)$ (and so $v \in N_A[H_y]$), and it is a solid edge because v is not an isolated vertex of H_y . If $u \notin H_y$, then by Lemma 5.1, $H_y \subseteq H_x$. So, $u \in N_A(H_y)$ since $uy \in E(G)$, and this contradicts u being isolated in H_x . \Box **Lemma 5.3** Let C be a cycle of G of length at least 4 with no solid chord. If C is not a hole then there exist three consecutive vertices x, y, u in C such that:

- $u \in A$, $x, y \in B$, $\{u\} \subseteq H_y \subseteq H_x$ and u is an isolated vertex of H_x , or - $u \in A'$, $x, y \in B'$, $\{u\} \subseteq H'_y \subseteq H'_x$ and u is an isolated vertex of H'_x . In particular ux is an optional edge of G and a chord of C.

Proof.

We may assume that C has a chord e for otherwise it is a hole. This chord cannot be a flat edge of G because a flat edge contains a vertex of I, so a vertex of degree 2, and it therefore cannot be a chord of any cycle. Hence, e is an optional edge of G. So, up to symmetry, we may assume that e = ux with $u \in A$ and $x \in B$. By definition of optional edges, u is an isolated vertex of $G[H_x]$.

Let u' and y be the two neighbors of u along C. If $u', y \in A \cup B$, then by Lemma 5.2, u'y is a solid chord of C, a contradiction. So, up to symmetry, $y \in A \cup B$ and $u' \in I$.

Suppose first that $y \in A$. Since $uy \in E(G)$ and u is isolated in H_x , we have that $y \in N_A(H_x)$. If follows that $xy \in E(G)$, and moreover, xy is a solid edge since $y \notin H_x$. Since x and y are both in C and C has no solid chord, C visits consecutively u', u, y and x. Let x' be the neighbor of x in $C \setminus y$. If $x' \in B$ then $H_x \cap H_{x'} \neq \emptyset$, and since y is complete to H_x , y has a neighbor in $H_{x'}$. It follows that yx' is an edge of G, the edge yx' is solid, and is therefore a solid chord of C, a contradiction. Hence $x' \in A$, and so since xx' is an edge, $x' \in N_A[H_x]$. If $x' \in H_x$, then yx' is a solid chord of C, and if $x' \in N(H_x)$, then (since H_x is a module of G[A]) x'u is a solid chord of C, in each case a contradiction.

Suppose now that $y \in B$. If $H_y \subseteq H_x$, then xy is an edge that is solid and hence is an edge of C, so the conclusion of the lemma holds. So we may assume by Lemma 5.1 that $H_x \subseteq H_y$. In particular, xy is an edge, and since it is solid, u, y and x are consecutive along C. Let v be the neighbor of xin $C \setminus y$. If $v \in B$, then $H_v \cap H_x \neq \emptyset$, so $H_v \cap H_y \neq \emptyset$, showing that yv is a solid chord of G, a contradiction. Hence, $v \in A$. We have $uv \notin E(G)$ for otherwise uv would be a solid chord of G. Hence, $v \in H_x$ since $vx \in E(G)$ and H_x is a module of G[A]. So, $v \in H_y$ (and hence $vy \in E(G)$) and v is an isolated vertex of H_y , for otherwise vy would be a solid chord of G.

Now, we have three consecutive vertices y, x, v in C such that: $v \in A$, $y, x \in B$, $\{v\} \subseteq H_x \subseteq H_y$ and v is an isolated vertex of H_y . So, the conclusion of the lemma is satisfied again with these three vertices. \Box

5.2 Blowups and holes

Let G be a twinless odd ℓ -template with an ℓ -partition (A, B, A', B', I, w, w')A blowup of G is any graph G^* that satisfies the following:

- (a) For every vertex u of G there is a clique K_u in G^* on $k_u \ge 1$ vertices u_1, \ldots, u_{k_u} such that $u_{k_u} = u$; for distinct vertices u, v of $G, K_u \cap K_v = \emptyset$ and $V(G^*) = \bigcup_{u \in V(G)} K_u$, so $V(G) \subseteq V(G^*)$.
- (b) For all vertices $u \in V(G)$ and all integers $1 \leq i \leq j \leq k_u$, in G^* $N[u_i] \subseteq N[u_j]$ (in particular, for all $u, v \in V(G)$, $G^*[K_u \cup K_v]$ is a half graph).
- (c) If u and v are non-adjacent vertices of G, then K_u is anticomplete to K_v (in particular $uv \notin E(G^*)$).
- (d) If uv is a solid edge of G, then K_u is complete to K_v (in particular $uv \in E(G^*)$).
- (e) If uv is a flat edge of G, then u is complete to K_v and v is complete to K_u (in particular $uv \in E(G^*)$).
- (f) If ux is an optional edge of G with $u \in A$ and $x \in B$ (resp. $u \in A'$ and $x \in B'$), then u is complete to K_x (in particular $uv \in E(G^*)$).
- (g) If ux and uy are optional edges of G with $u \in A$, $x, y \in B$ and $H_y \subsetneq H_x$ (resp. $u \in A', x, y \in B'$ and $H'_y \subsetneq H'_x$), then every vertex of K_u with a neighbor in K_y is complete to K_x .
- (h) w (resp. w') is a universal vertex of $G^*[\bigcup_{u \in A \cup B} K_u]$ (resp. $G^*[\bigcup_{u \in A' \cup B'} K_u]$).

Observe that $G = G^*[V(G)]$ follows clearly from the definition, so G is an induced subgraph of G^* . For every vertex u of G, the clique K_u is called a blown up clique, more specifically the clique blown up from u.

Note that to define the blowup of a graph, it is first needed to fix an ℓ -partition of it. Also, it should be stressed that the blowup is defined only for twinless graphs. Hence, in condition (g) of the definition, since G is twinless, when $x \neq y$, $H_y \subsetneq H_x$ is equivalent to $H_y \subseteq H_x$ because $H_x = H_y$ would imply that x and y are twins.

Lemma 5.4 A hole C in a blowup of a twinless odd ℓ -template contains at most one vertex in each blown up clique.

Proof. Since a hole is triangle-free, C intersects any clique in at most two vertices. So suppose for a contradiction that some blown up clique K_v contains two vertices x and y of C. Let x' be the neighbor of x in $C \setminus y$ and y' be the neighbor of y in $C \setminus x$. Since by condition (b) of the definition of the blowup we have that in G^* , $N[x] \subseteq N[y]$ or $N[y] \subseteq N[x]$, one of xyx' or xyy' is a triangle of C, a contradiction.

Lemma 5.5 In a blowup G^* of a twinless odd ℓ -template G, every hole has length $2\ell + 1$.

Proof. Let C^* be a hole in G^* . By Lemma 5.4, it contains at most one vertex in each blown up clique. Let C be the subgraph of G that is induced by all vertices v such that some vertex of C^* is in K_v . By Lemma 5.4, $|V(C^*)| = |V(C)|$. By the definition of blowup (specifically conditions (c) and (d)), C^* is isomorphic to some graph obtained from C by removing optional or flat edges of G. Hence, C is a cycle of G with no solid chord. If C is a hole of G, then since it has the same length as C^* , by Lemma 4.11, C^* has length $2\ell + 1$. Hence, we may assume that C has chords, so by Lemma 5.3, without loss of generality, C contains three consecutive vertices x, y, u such that: $u \in A, x, y \in B$, $\{u\} \subsetneq H_y \subsetneq H_x$ and u is an isolated vertex of H_x . Note that it follows that both ux and uy are optional edges of G. Because of C^* , the vertex u_i of $K_u \cap V(C^*)$ has a neighbor in K_y . So, by condition (g) of blowups, u_i is complete to K_x . Hence, C^* has a chord, a contradiction.

5.3 Preblowup

Checking that a graph is the blowup of a template is tedious. Here we provide a simpler notion and prove it is in some sense equivalent.

A preblowup of an odd ℓ -template G with an ℓ -partition (A, B, A', B', I, w, w') is any graph G^* obtained from G as follows. Every vertex u of $A \cup A' \cup I$ is replaced by a clique K_u on $k_u \geq 1$ vertices such that $u \in K_u$. We denote by A^* the set $\bigcup_{u \in A} K_u$ and use a similar notation A'^* and I^* . The set B (resp. B') is replaced by a set B^* (resp. B'^*) of vertices such that $B \subseteq B^*$ (resp. $B' \subseteq B'^*$). So, $V(G^*) = A^* \cup B^* \cup A'^* \cup B'^* \cup I^*$. The sets $A^*, B^*, A'^*, B'^*, I^*$ are disjoint. Vertices of G are adjacent in G^* if and only if they are adjacent in G, so G is an induced subgraph of G^* . Finally, we require that the following conditions hold (throughout N refers to the neighborhood in G^*):

- (a) For all $u \in A$, $N(K_u) \subseteq A^* \cup B^* \cup K_{u^+}$ where u^+ is the neighbor of u in I and:
 - 1. For every $u^* \in K_u$, $N_A(u^*) = N_A[u]$.
 - 2. Every vertex of K_u has a neighbor in K_{u^+} .
- (b) $N(B^*) \subseteq A^*$ and:
 - 1. If $w \in B$, then there exists $w^* \in B^*$ that is complete to A^* .
 - 2. If $u^* \in B^*$, then there exist non-adjacent $a, b \in A$ such that u^* has neighbors in both K_a and K_b .
- (i) For all $u \in I$, $N(K_u) \subseteq K_a \cup K_b$ where a and b are the neighbors of u in G, and:
 - 1. Every vertex $u^* \in K_u$ has at least one neighbor in each of K_a and K_b .

Conditions (a') and (b') analogous to (a) and (b) hold for A' and B'.

Recall that to blowup (resp. preblowup) a template, one needs to first fix an ℓ -partition. If this partition is proper, the blowup (resp. preblowup) is *proper*. Recall that by Lemma 4.16, a proper ℓ -partition (A, B, A', B', I, w, w') exists for every twinless odd ℓ -template G (but this remark will be used only in the next section, so far we just assume the ℓ -partition we work with is proper).

When G^* is a preblowup of a template G, the *domination score* of G w.r.t. G^* is (where N refers to the neighborhood in G^*):

$$s(G, G^*) = \sum_{x \in A \cup A' \cup I} |\{x^* \in K_x : N[x^*] \subseteq N[x]\}|$$

Observe that the blowup is defined only for twinless templates while the preblowup is defined for any template. It is straightforward to check that a blowup is a particular preblowup. The following is a converse of this statement.

Lemma 5.6 Let $\ell \geq 3$ and let G^* be a proper preblowup of an odd ℓ -template with $k \geq 3$ principal paths. If $G^* \in C_{2\ell+1}$, then G^* is a proper blowup of a twinless odd ℓ -template G with k principal paths (in particular, G is an induced subgraph of G^*). *Proof.* Among all the induced subgraphs of G^* that are odd ℓ -templates and for which G^* is a proper preblowup, we suppose that G is one that maximizes $s(G, G^*)$. We denote by (A, B, A', B', I, w, w') the proper ℓ -partition of Gthat is used for its preblowup and by $(A^*, B^*, A'^*, B'^*, I^*)$ the corresponding partition of the vertices of G^* .

(1) There exist vertices w^* and w'^* that are complete to respectively $A^* \setminus \{w^*\}$ and $A'^* \setminus \{w'^*\}$, and such that either $w^* \in B^*$ and $w'^* \in A'^*$, or $w^* \in A^*$ and $w'^* \in B'^*$.

If $w \in A$, then from the definition of w (see Lemma 4.4), the definition of A^* and condition (a1), it follows that $w^* = w$ is complete to $A^* \setminus \{w^*\}$. If $w \in B$, by condition (b1) there exists $w^* \in B^*$ that is complete to A^* .

The statement about w'^* holds by symmetry. The last statement comes from the fact that by Lemma 4.4 exactly one of w, w' is in $A \cup A'$, and the other one is in $B \cup B'$. This proves (1).

(2) For every principal path $P_u = u \dots u'$ of G and $u^* \in K_u$, there exists in G^* a path P_{u^*} of length $\ell - 1$ from u^* to some $u'^* \in K_{u'}$ whose interior is in $\bigcup_{x \in I \cap V(P_u)} K_x$. Moreover, the interior of P_{u^*} is anticomplete to $V(G^*) \setminus \bigcup_{v \in V(P_u)} K_v$.

The existence of a path from u^* to some $u'^* \in K_{u'}$ whose interior is in $\bigcup_{x \in I \cap V(P)} K_x$ follows from conditions (a), (i), (i1), and (a2) of preblowup. Its length is $\ell - 1$ by condition (c) of templates. The statement about its interior follows from conditions (a), (b) and (i) of preblowup. This proves (2).

(3) For all $u, v \in A$ such that $uv \notin E(G)$, K_u is anticomplete to K_v . A similar statement holds for A'.

Suppose that there exists $u^* \in K_u$ and $v^* \in K_v$ such that $u^*v^* \in E(G^*)$. By condition (a1) of preblowup, $u \neq u^*$ and $v \neq v^*$. Let $P_u = u \dots u'$ and $P_v = v \dots v'$ be principal paths. Denote by u^+ the neighbor of u in P_u and by v^+ the neighbor of v in P_v . By property ((i)) of a template, $u'v' \in E(G)$. Hence $uP_u u'v' P_v vv^* u^* u$ is a cycle C. By conditions (a) and (a1) of preblowup, the only possible chords in C are u^+u^* and v^+v^* . Without loss of generality, we may assume that $u^*u^+ \in E(G^*)$ for otherwise C is a hole of length $2\ell + 2$, a contradiction.

Let P_{v^*} be a path of length $\ell - 1$ from v^* to v'^* as defined in (2). Since $v'^* \in K_{v'}$ and by (a1) applied to $A', v'^*u' \in E(G^*)$ and $v^*P_{v^*}v'^*u'P_uu^+u^*v^*$ is a hole of length 2ℓ , a contradiction.

The result for A' holds symmetrically. This proves (3).

(4) For all $u, v \in A$ such that $uv \in E(G)$, K_u is complete to K_v . A similar statement holds for A'.

Suppose that there exists $u^* \in K_u$ and $v^* \in K_v$ such that $u^*v^* \notin E(G^*)$. Let $P_{u^*} = u^* \dots u'^*$ and $P_{v^*} = v^* \dots v'^*$ be defined as in (2). Observe that $u'^* \in K_{u'}$ and $v'^* \in K_{v'}$. Furthermore $u'v' \notin E(G)$ by property ((i)) of templates. Hence, by (3), $u'^*v'^* \notin E(G^*)$.

We claim that there exists a vertex $a \in (A \cup B) \setminus \{u, v\}$ that is adjacent to both u^* and v^* . If $w^* \neq u, v$, then by (1) and condition (a2), we may choose $a = w^*$. Otherwise, up to symmetry, $w^* = u$. Since the ℓ -partition of G is proper, by Lemma 4.17, A contains a universal vertex x distinct from $w^* = u$. If $x \neq v$, we set a = x. If x = v, then both u and v are universal vertices of G[A] and we may choose for a any vertex of $A \setminus \{u, v\}$. This proves our claim.

Now, $au^*P_{u^*}u'^*v'^*P_{v^*}v^*a$ is a hole of length $2\ell + 2$, a contradiction. The result for A' holds symmetrically. This proves (4).

(5) For all $u \in I$ and $u_1, u_2 \in K_u$, either $N[u_1] \subseteq N[u_2]$ or $N[u_2] \subseteq N[u_1]$.

Otherwise, there exists $x_1^* \in N[u_1] \setminus N[u_2]$ and $x_2^* \in N[u_2] \setminus N[u_1]$. Note that $x_1^*x_2^* \notin E(G^*)$ for otherwise, $\{x_1^*, x_2^*, u_1, u_2\}$ induces a C_4 . It follows that x_1^* and x_2^* belong respectively to distinct cliques K_{x_1} and K_{x_2} , where x_1 and x_2 are the two neighbors of u along some principal path $P = v \dots v'$ of G. Because of x_1^*, x_2^* and condition (i1) of preblowup, there exists a path P^* of length ℓ from some $v^* \in K_v$ to some $v'^* \in K_{v'}$ whose interior is in $\bigcup_{x \in I \cap V(P)} K_x$.

Let $q \neq v$ be a vertex of A and $Q = q \dots q'$ be a principal path of G, and suppose up to symmetry that $qv \notin E(G)$. Now, by conditions (i) and (a1) of preblowup and (1), P^* , Q and w^* form a hole of length $2\ell + 2$. This proves (5).

(6) For all $u \in I$ and $u^* \in K_u$, $N[u^*] \subseteq N[u]$.

Otherwise, by (5), there exists a vertex $u^* \in K_u$ such that $N[u] \subsetneq N[u^*]$. Hence $(V(G) \setminus \{u\}) \cup \{u^*\}$ induces a subgraph G_0 of G^* and it is easy to verify that G^* is a preblowup of G_0 . This contradicts to the maximality of $s(G, G^*)$. This proves (6).

By (5), for every $u \in I$, the clique K_u can be linearly ordered by the inclusion of the neighborhoods as u_1, \ldots, u_{k_u} with $u = u_{k_u}$ by (6) (so, for $1 \leq i \leq j \leq k_u$, $N[u_i] \subseteq N[u_j]$). From condition (i) of the preblowup it also follows that, in G^* , u is complete to the cliques associated to its two

neighbors in G.

(7) For every $u \in A$ and $u_1, u_2 \in K_u$, either $N[u_1] \subseteq N[u_2]$ or $N[u_2] \subseteq N[u_1]$. A similar statement holds for A'.

Otherwise, there exist $x_1 \in N[u_1] \setminus N[u_2]$ and $x_2 \in N[u_2] \setminus N[u_1]$. Note that $x_1x_2 \notin E(G^*)$ for otherwise, $\{x_1, x_2, u_1, u_2\}$ induces a C_4 .

Observe first that by (3) and (4), $N_{A^*}[u_1] = N_{A^*}[u_2]$. Hence by (a) of preblowup, $x_1, x_2 \in B^* \cup K_{u^+}$ where u^+ is the neighbor of u in the principal path that contains u. Without loss of generality and since K_{u^+} is a clique, $x_1 \in B^*$.

By condition (b2), there exist non-adjacent $a, b \in A$ such that x_1 has neighbors $a^* \in K_a$ and $b^* \in K_b$, and by (3) $a^*b^* \notin E(G^*)$. Note that $a^*, b^* \neq$ u_2 because $u_2x_1 \notin E(G^*)$. If u_2 is complete to $\{a^*, b^*\}$, then $\{u_2, a^*, x_1, b^*\}$ induces a C_4 , a contradiction. So, up to symmetry $u_2a^* \notin E(G)$. So, $a^* \notin K_u$ and by (4) and (3), $a^*u_1 \notin E(G^*)$. Observe that $x_2a^* \notin E(G^*)$ for otherwise $\{a^*, x_1, u_1, u_2, x_2\}$ induces a C_5 .

Suppose that $x_2 \in B^*$. As above, we can show that x_2 has a neighbor $c^* \in A^*$ that is anticomplete to $\{u_1, u_2, x_1\}$. Note that $a^*c^* \notin E(G^*)$ for otherwise $\{x_1, a^*, c^*, x_2, u_2, u_1\}$ induces a C_6 . Let $P_{a^*} = a^* \dots a'^*$ and $P_{c^*} = c^* \dots c'^*$ be defined as in (2).

By (3) and (4) and since $a^*c^* \notin E(G^*)$, $a'^*c'^* \in E(G^*)$. So, by conditions (a), (b) and (i), $u_1x_1a^*P_{a^*}a'^*c'^*P_{c^*}c^*x_2u_2u_1$ is a hole of length $2\ell + 4$, a contradiction.

So $x_2 \in K_{u^+}$. Hence by condition (i1) of preblowup, there exists a path Q of length $\ell - 2$ from x_2 to some $u'^* \in K_{u'}$. Now $x_2Qu'^*a'^*P_{a^*}a^*x_1u_1u_2x_2$ is a hole of length $2\ell + 2$, a contradiction.

The result for A' holds symmetrically. This proves (7).

(8) For all $u \in A$ and $u^* \in K_u$, $N[u^*] \subseteq N[u]$. A similar statement holds for A'.

Otherwise, there exists a vertex $u^* \in K_u$ such that $N[u] \subsetneq N[u^*]$. Hence, $(V(G) \setminus \{u\}) \cup \{u^*\}$ induces a subgraph G_0 of G^* and it is easy to verify that G^* is a preblowup of G_0 (that is a template by Lemma 4.14 and whose partition is proper by (3) and (4)). This contradicts the maximality of $s(G, G^*)$. The result for A' holds symmetrically. This proves (8).

By (7), for every $u \in A \cup A'$, the clique K_u can be linearly ordered by the inclusion of the neighborhoods as u_1, \ldots, u_{k_u} , and by (8) $u_{k_u} = u$ (so, for $1 \leq i \leq j \leq k_u$, $N[u_i] \subseteq N[u_j]$).

(9) If xy is an edge of $G[B^*]$, then either $N_{A^*}(x) \subseteq N_{A^*}(y)$ or $N_{A^*}(y) \subseteq$

 $N_{A^*}(x).$

Otherwise, there exists $u^* \in N_{A^*}(x) \setminus N_{A^*}(y)$ and $v^* \in N_{A^*}(y) \setminus N_{A^*}(x)$. Note that $u^*v^* \notin E$ for otherwise $\{u^*, x, y, v^*\}$ induces a C_4 . So, for some $u, v \in A$, we have $u^* \in K_u$ and $v^* \in K_v$. Hence, by (4), $uv \notin E(G)$. Let $P_{u^*} =$ $u^* \dots u'^*$ and $P_{v^*} = v^* \dots v'^*$ be defined as in (2). So, $xu^*P_{u^*}u'^*v'^*P_{v^*}v^*yx$ form a hole of length $2\ell + 2$, a contradiction. This proves (9).

(10) For every $x \in B^*$, there exist non-adjacent $u, v \in A$ such that $xu, xv \in E(G^*)$.

This follows from condition (b2) of preblowup and from (8). This proves (10).

Two vertices x, y in B^* are equivalent if $N_A(x) = N_A(y)$.

(11) If x and y are equivalent vertices of B^* , then $xy \in E(G^*)$.

If $xy \notin E(G^*)$, then x, y and two of their neighbors provided by (10) induce a C_4 . This proves (11).

Vertices of B^* are partitioned into equivalence classes. By (11), each equivalence class is a clique X, and by (9), vertices of X can be linearly ordered according to the inclusion of neighborhoods in A^* . In each such a clique X we choose a vertex x maximal for the order and call B_1 the set of these maximal vertices. For every $x \in B_1$, we denote by K_x the clique of B^* of all vertices equivalent to x. Observe that if $w^* \in B$, then w^* is a maximal vertex of its clique. Hence, we can set $w^* \in B_1$.

So, for every $u \in B_1$, the clique K_u can be linearly ordered by the inclusion of the neighborhod in A^* as u_1, \ldots, u_{k_u} with $u = u_{k_u}$ (so, for $1 \le i \le j \le k_u, N_{A^*}(u_i) \subseteq N_{A^*}(u_j)$).

Statements similar to (9), (10), (11) hold for $B^{\prime*}$ and we define B_1^{\prime} as well.

We set $G_1 = G^*[A \cup B_1 \cup A' \cup B'_1 \cup I]$ and claim that (A, B_1, A', B'_1, I) is an ℓ -pretemplate partition of G_1 . Since $G_1[A \cup I \cup A']$ is exactly $G[A \cup I \cup A']$, conditions (c), (d), (e) and (f) hold. Adding the fact that $N_{G_1}(B_1) \subseteq A^* \cap$ $V(G_1) = A$ by condition (b) of preblowup, condition (a) for a pretemplate holds and symmetrically also condition (b). Now condition (g) holds because w^* and w'^* are complete to respectively $A \cup B_1$ and $A' \cup B'_1$. By (10), the last two conditions for a pretemplate are fulfilled by (A, B_1, A', B'_1, I) . Hence, by Lemma 4.14, G_1 is a an odd ℓ -template. It is twinless by Lemma 4.15. We also notice that by construction w^* and w'^* belong to G_1 . Furthermore, by (1), w^* (respectively w'^*) is complete to $A \setminus \{w^*\}$ (respectively $A' \setminus \{w'^*\}$). From the definition of a template it is easy to conclude that w^* (respectively $w^{\prime*}$) is universal in $G_1[A \cup B_1]$ (respectively $G_1[A^{\prime} \cup B_1^{\prime}]$). Hence $(A, B_1, A^{\prime}, B_1^{\prime}, I, w^*, w^{\prime*})$ is a proper ℓ -partition of G_1 .

We now prove that G^* is a proper blowup of G_1 .

By the definition of a preblowup and by (11), for all $u \in V(G_1)$, K_u is a clique and $V(G^*) = \bigcup_{u \in V(G_1)} K_u$

(12) If
$$u, v \in V(G_1)$$
 and $uv \notin E(G_1)$, then K_u is anticomplete to K_v .

Suppose $u, v \in V(G_1)$ and $uv \notin E(G_1)$. If $u \in I$ or $v \in I$, the conclusion follows directly from condition (i) of preblowup. So we may assume up to symmetry that $u \in A \cup B_1$. By conditions (a) and (b) of preblowup, we may assume $v \in A \cup B_1$. If $u, v \in A$, then the result follows from (3), so we may assume that $v \in B_1$.

Now suppose for a contradiction that there exist $u^* \in K_u$ and $v^* \in K_v$ such that $u^*v^* \in E(G_1)$. By the choice of vertices in B_1 , for all $v^* \in K_v$, $N[v^*] \subseteq N[v]$. So $u^*v \in E(G_1)$. For the same reason or by (8), for all $u^* \in K_u$, $N[u^*] \subseteq N[u]$. Hence $uv \in E(G_1)$, a contradiction. This proves (12).

(13) If uv is a solid edge of G_1 then K_u is complete to K_v .

Otherwise, let $u^* \in K_u$ and $v^* \in K_v$ such that $u^*v^* \notin E(G)$. Since uv is a solid edge, up to symmetry, $u, v \in A$ or $u, v \in B_1$ or $u \in A$, $v \in B_1$ and in this last case u is not an isolated vertex of $G[H_v]$.

By (4) the case where u and v are in A cannot happen. Assume then that $v \in B_1$. By Lemma 4.6, there exist $a, b \in H_v$ (and hence in A) that are not adjacent. Assume that u is also in B_1 . Since u and v are adjacent, by (9) we may assume without loss of generality that $H_v \subseteq H_u$ and so aand b belong to H_u too. Then, by the definition of K_u and K_v , we get a C_4 induced by $\{u^*, v^*, a, b\}$, a contradiction. So u should be in A, and to avoid a C_4 induced by $\{u^*, v^*, a, b\}$, u^* should be non-adjacent to at least one of a and b, say a. In particular, $a \neq u$. Then, by (4), $ua \notin E(G_1)$. So u does not belong to $N(H_v)$ and since uv is an edge of G_1 , we get that $u \in H_v$. Since uv is solid, u has at least one neighbor in H_v , and it is not adjacent to $a \in H_v$. Hence, as H_v is anticonnected, there exist non-adjacent vertices $c, d \in H_v$ such that $uc \notin E(G_1)$ and $ud \in E(G_1)$. Now $u^*P_{u^*}u'^*c'P_ccv^*du^*$ is a hole of length $2\ell + 2$, a contradiction again.

This proves (13).

(14) For all $u \in V(G_1)$ and $1 \le i \le j \le k_u$, $N[u_i] \subseteq N[u_j]$.

The result follows from how vertices are ordered after the proof of (5) (vertices in I), (7) (vertices in A or A') and (11) (vertices in B_1 or B'_1). This proves (14).

(15) If uv is a flat edge of G_1 , then u is complete to K_v and v is complete to K_u .

By definition of a flat edge, either u and v are in I or one is in I and the other is in A or in A'. The result follows from (6), (8), and conditions (a2) (applied to A or A') and (i1) of the preblowup. This proves (15).

(16) If ux is an optional edge of G_1 with $u \in A$ and $x \in B_1$ (resp. $u \in A'$ and $x \in B'_1$), then u is complete to K_x .

The result follows from the definition of K_x when $x \in B_1$. This proves (16).

(17) If ux and uy are optional edges with $u \in A$, $x, y \in B_1$ and $H_y \subsetneq H_x$ (resp. $u \in A'$, $x, y \in B'_1$ and $H'_y \subsetneq H'_x$), then every vertex of K_u with a neighbor in K_y is complete to K_x .

Otherwise, let u^* be a vertex in K_u that has a neighbor y^* in K_y and a non-neighbor x^* in K_x . Since H_x and H_y are not disjoint, xy is a solid edge of G_1 and by (13), $x^*y^* \in E(G_1)$.

Since x and y are not equivalent, there exists a vertex a such that $a \in N_A(y) \setminus N_A(x)$ or $a \in N_A(x) \setminus N_A(y)$. In the first case, by definition of a template, $a \in A \setminus N_A[H_x]$. Then since $H_y \subsetneq H_x$ and H_x is a module of A we get that a is anticomplete to H_x and hence to H_y . So $a \notin N_A(y)$, a contradiction; we may then conclude that $a \in N_A(x) \setminus N_A(y)$

By definition of the cliques in B, $x^*a \in E(G^*)$ and $y^*a \notin E(G^*)$. Therefore, to avoid a C_4 induced by $\{x^*, y^*, u^*, a\}$, it should be that $u^*a \notin E(G^*)$.

Now $aP_aa'u'^*P_{u^*}u^*y^*x^*a$ is a hole of length $2\ell + 2$ a contradiction. This proves (17).

(18) w^* (resp. w'^*) is a universal vertex of $G^*[\bigcup_{u \in A \cup B_1} K_u]$ (resp. $G^*[\bigcup_{u \in A' \cup B'_1} K_u]$).

By (1), w^* is complete to $A^* \setminus \{w^*\}$ and so to $\bigcup_{u \in A} K_u \setminus \{w^*\}$. Furthermore, from the definition of G_1 we know that w^* is complete to $B_1 \setminus \{w^*\}$. If $w^* \in B_1$, since all edges between vertices in B_1 are solid, by (13), w^* is complete to $B^* \setminus \{w^*\}$. Similarly, if $w^* \in A$, by (13) and (16), we get that w^* is complete to B^* . In both cases w^* is a universal vertex of $G^*[\bigcup_{u \in A \cup B} K_u]$. The proof for w'^* is symmetric. This proves (18).

From all the claims above, G^* satisfies all conditions to be a proper blowup of G_1 .
6 Graphs in $C_{2\ell+1}$ that contain a pyramid

The goal of this section is to prove the following.

Lemma 6.1 Let $\ell \geq 3$ be an integer. If G is a graph in $C_{2\ell+1}$ and G contains a pyramid, then one of the following holds:

- (a) G is a proper blowup of a twinless odd ℓ -template;
- (b) G has a universal vertex;
- (c) G has a clique cutset.

The rest of this section is devoted to the proof of Lemma 6.1. So from here on $\ell \geq 3$ is an integer and G is a graph in $\mathcal{C}_{2\ell+1}$ that contains a pyramid Π . By Lemma 3.10, the three paths of Π have length ℓ . By Lemma 4.5, Π is an odd ℓ -template. Hence, we may define an integer k and a sequence F_0, F_1, F_2 of induced subgraphs of G as follows.

- k is the maximum integer such that G contains an odd l-template with k principal paths. Observe that by Lemma 4.15, G in fact contains a twinless template with k principal paths, because twins can be eliminated from templates by deleting hyperedges with equal vertex-set while there are some.
- In G, pick a proper blowup F_1 of a twinless odd ℓ -template F_0 with k principal paths. Note that F_0 exists and the proper ℓ -partition needed for the proper blowup exists by Lemma 4.16.
- Suppose that F_0 and F_1 are chosen subject to the maximality of the vertex-set of F_1 (in the sense of inclusion). Note that possibly F_0 is not a maximal template in the sense of inclusion, it can be that a smaller template leads to a bigger blowup (but F_0 has k principal paths).
- F_2 is obtained from F_1 by adding all vertices of $G \setminus F_1$ that are complete to F_1 .

Lemma 6.2 $V(F_2) \setminus V(F_1)$ is a (possibly empty) clique that is complete to F_1 .

Proof. Otherwise, G contains a C_4 .

We now introduce some notation. We denote by (A, B, A', B', I, w, w')the proper ℓ -partition that is used to blow up F_0 . When u is a vertex of F_0 , we denote by K_u the clique of F_1 that is blown up from u. We set $A^* = \bigcup_{u \in A} K_u$. We use a similar notation B^* , A'^* , B'^* and I^* .

6.1 Technical lemmas

We now prove lemmas that sum up several structural properties of G.

Lemma 6.3 If $u \in A \cup A' \cup I \cup \{w, w'\}$ and $v \in N_{V(F_0)}(u)$, then u is complete to K_v .

Proof. We prove this lemma using the conditions from the definition of blowups. If $u \in \{w, w'\}$, then the result follows from condition (h). If $u \in A \cup A'$, then the conclusion follows from conditions (d), (e) and (f). If $u \in I$, then the conclusion follows from condition (e).

Very often, Lemma 6.3 will be used in the following way. Suppose there exists a principal path $P = u \dots u'$ of F_0 . Suppose there exists a vertex x of P and $x^* \in K_x$. Then by Lemma 6.3 and condition (c) of blowups, $\{x^*\} \cup (V(P) \setminus \{x\})$ induces a path of F_1 . If $y \neq x$ is a vertex of P and $y^* \in K_y$, then $\{x^*, y^*\} \cup (V(P) \setminus \{x, y\})$ might fail to induce a path of F_1 , because it is possible that $xy \in E(G)$ while $x^*y^* \notin E(G)$. But under the assumption that $x^*y^* \in E(G)$ or $xy \notin E(G)$, we do have that $\{x^*, y^*\} \cup (V(P) \setminus \{x, y\})$ induces a path of F_1 . Several variant of this situation will appear soon and we will simply justify them by referring to Lemma 6.3.

When u is a vertex in A, we denote by P_u the unique principal path of F_0 that contains u. Its end in A' is then denoted by u'. We denote by u^+ the neighbor of u in P_u . We denote by u^{++} the neighbor of u^+ in $P_u \setminus u$. Note that $u^+ \in I$ and $u^{++} \in I \cup A'$ ($u^{++} \in A'$ if and only if $\ell = 3$).

For any distinct $u, v \in A$, from the definition of templates, exactly one of $V(P_u) \cup V(P_v) \cup \{w\}$ or $V(P_u) \cup V(P_v) \cup \{w'\}$ induces a hole that is denoted by $C_{u,v}$. Such a hole is called a *principal hole*.

Note that there are two kinds of principal holes: those that contain w, and those that contain w'. Recall that by Lemma 4.11, every hole of a template contains two principal paths plus an extra vertex, but it may fail to be a principal hole (because it may fail to contain w or w'). Though we do not use this information formally, it is worth noting that by Lemma 6.3, when C is a principal hole, $\bigcup_{v \in V(C)} K_v$ induces a ring. But when C is a non-principal hole, it may happen that $\bigcup_{v \in V(C)} K_v$ does not induce a ring (because there might be in C an optional edge uv with $u \in A$ and $v \in B$, and after the blowup process, there might be no vertex in K_v that is complete to K_u).

Lemma 6.4 If $u \in V(F_0)$ and $u^* \in K_u$, then u^* has two neighbors in $V(F_0) \setminus K_u$ that are not adjacent.

Proof. If $u \in I$, then let P be the principal path that contains u. By Lemma 6.3, u^* is adjacent to the two neighbors of u in P.

If $u \in A \cup A'$, say $u \in A$ up to symmetry, then we claim that u has a neighbor z in $A \cup B$. This is clear if u is not isolated in A and otherwise we set z = w. By Lemma 6.3, z and u^+ are non-adjacent neighbors of u^* .

If $u \in B$, then by the definition of a template, H_u contains two non adjacent vertices a and b that are neighbors of u. By Lemma 6.3, a and b are both adjacent to u^* .

Lemma 6.5 If uv is an edge of $F_0[A \cup A' \cup I \cup \{w, w'\}]$, then some principal hole of F_0 goes through uv.

Proof. If at least one of u, v is in I then uv is an edge of a principal path and we know that this principal path belongs to a principal hole. Else, since $A \cup \{w\}$ is anticomplete to $A' \cup \{w'\}$, up to symmetry both u and v are in A or $u = w \in B$ and $v \in A$.

If $u, v \in A$ then $C_{u,v}$ is a principal hole containing uv.

If $u = w \in B$ and $v \in A$: since w is in B, G[A] has no universal vertex and there exists $a \in A$ which is not adjacent to v. Now w, P_v, P_a form a principal hole containing the edge uv.

Lemma 6.6 If K is a clique of F_0 , $K^* = \bigcup_{v \in K} K_v$ and D is a connected induced subgraph of $G \setminus F_2$ such that $N_{V(F_1)}(D) \subseteq K^*$, then $N_{V(F_1)}(D)$ is a clique.

Proof. For suppose not. This means that there exists $u^*, v^* \in K^*$ and $x_u, x_v \in D$ such that $u^*v^* \notin E(G)$ and $x_uu^*, x_vv^* \in E(G)$ (possibly $x_u = x_v$). Since D is connected, there exists a path P in D from x_u to x_v . Suppose that u^*, x_u, v^*, x_v and P are chosen subject to the minimality of P. It follows that $u^*x_uPx_vv^*$ is a path, and recall that by assumption its interior is anticomplete to $F_1 \setminus K^*$.

Since $u^*v^* \notin E(G)$, u^* and v^* are in different blown-up cliques. Denote by K_u and K_v the blown-up cliques such that $u^* \in K_u$ and $v^* \in K_v$. By hypothesis, $uv \in K$ and so $uv \in E(G)$. Since $u^*v^* \notin E(G)$, by condition (d) of blowups, uv is not a solid edge of G.

If uv is a flat edge of F_0 , then by Lemma 6.5 a principal hole C goes through uv. Note that apart from u and v, no vertex of C is in K since K is a clique. By Lemma 6.3, in G, $(\{u^*, v^*\}) \cup V(C)) \setminus \{u, v\}$ induces a path Q of length 2ℓ . So P and Q form a hole of length at least $2\ell + 2$, a contradiction. If uv is an optional edge of F_0 , say with $u \in A$ and $v \in B$, then $u \in H_v$, and there exists a in H_v such that $au \notin E(F_0)$. Therefore, P_u , P_a and v form a hole C^* . By condition (f) of blowups (if va is optional), or by condition (d) (if va is solid), a is complete to K_v . By Lemma 6.3 it follows that $(\{u^*, v^*\}) \cup V(C^*)) \setminus \{u, v\}$ induces a path Q of length 2ℓ . So P and Qform a hole of length at least $2\ell + 2$, a contradiction again. \Box

When C is a hole of G, a vertex v of $V(G) \setminus V(C)$ is minor w.r.t. C if $N_{V(C)}(v)$ is included in a 3-vertex path of C. A vertex of $V(G) \setminus V(C)$ that is not minor w.r.t. C is major w.r.t. C.

Lemma 6.7 If $x \in V(G) \setminus V(F_2)$ and C is a principal hole of F_0 , then x is minor w.r.t. C.

Proof. Suppose up to symmetry that $w \in V(C)$ and suppose $C = C_{u,v}$. If x is major w.r.t. C, then C and x form a theta or a wheel that is not a twin-wheel. So by Lemma 3.10, x and C form a universal wheel. Let $P_t = t \dots t'$ be a principal path where $t \neq u, v$. If t is complete to $\{u, v\}$, then $xt \in E(G)$ for otherwise $\{t, u, v, x\}$ induces a C_4 . Hence x has at least 4 neighbors in $C_{u,t}$, so by Lemma 3.10, x is complete to P_t . If t is not complete to $\{u, v\}$, say $tu \notin E(G)$, then x again has at least 4 neighbors in $C_{u,t}$ because $w \in V(C_{u,t})$, so again x is complete to P_t .

We proved that x is complete to all principal paths, so to $I \cup A \cup A'$. Let $y \in B \cup B'$. By definition of a template y has two neighbors a and b, both in A or both in A', that are non-adjacent. Therefore a, b, y and x form a C_4 , unless x is adjacent to y. This proves that x is complete to $B \cup B'$, and so to $V(F_0)$.

Let z be a vertex of F_0 and $z^* \in K_z$. By Lemma 6.4, there exists $a, b \in V(F_0)$ such that $z^*a, z^*b \in E(G)$ and $ab \notin E(G)$, so since there is no C_4 in G it should be that $xz^* \in E(G)$. This proves that x is complete to F_1 . Hence, $x \in V(F_2)$, a contradiction.

Lemma 6.8 Let a and b be two non-adjacent vertices of some principal hole C of F_0 . If some vertex x of $V(G) \setminus V(F_2)$ has neighbors in both K_a and K_b , then a and b have a common neighbor c in C, x is adjacent to c, and x is anticomplete to every K_d such that $d \in V(C) \setminus \{a, b, c\}$.

Proof. Let $a^* \in K_a$ and $b^* \in K_b$ be two neighbors of x. Since $ab \notin E(G)$, by Lemma 6.3, $\{a^*, b^*\} \cup V(C) \setminus \{a, b\}$ induces a hole C^* . Since x is adjacent to a^* and b^* , by Lemma 3.10, x has another neighbor c in C^* (and in fact in

C since $c \neq a^*, b^*$). If c is not adjacent to a^* and b^* , then x is major w.r.t. C^* , so by Lemma 3.10, C^* and x form a universal wheel. It follows that x is major w.r.t. C, a contradiction to Lemma 6.7.

We proved that a and b have a common neighbor c in C and that x is adjacent to c. Suppose for a contradiction that x has a neighbor $d^* \in K_d$ where $d \in V(C) \setminus \{a, b, c\}$. By the same argument as above, since x has neighbors in K_d and K_c , c and d must have a common neighbor in C, and this common neighbor must be a or b, say a up to symmetry. So, x has neighbors in K_d and K_b while b and d have no common neighbors in C, so we may reach a contradiction as above. \Box

6.2 Connecting vertices of F_1

We here explain how lemmas of Subsection 4.4 are extended from F_0 to F_1 .

Lemma 6.9 If $u^* \in A^* \cup B^*$ and $v^* \in A'^* \cup B'^*$, then there exists in F_1 a path P^* of length $\ell - 1$, ℓ or $\ell + 1$ from u^* to v^* that contains the interior of a principal path.

More specifically:

- If $u^* \in A^*$ and $v^* \in A'^*$, then P^* has length $\ell 1$ or ℓ .
- If $u^* \in A^*$ and $v^* \in B'^*$, or if $u^* \in B^*$ and $v^* \in A'^*$, then P^* has length ℓ or $\ell + 1$.
- If $u^* \in B^*$ and $v^* \in B'^*$, then P^* has length $\ell + 1$.

Proof. Let u and v be such that $u^* \in K_u$ and $v^* \in K_v$. Let P be a path in F_0 like in Lemma 4.13 from u to v (so P contains the interior of some principal path Q). By Lemma 6.3, $\{u^*, v^*\} \cup V(P) \setminus \{u, v\}$ induces a path of the same length as P that contains the interior of Q.

Lemma 6.10 If $u^* \in B^*$ and $v^* \in B'^*$, then there exist in G two paths P^* and Q^* from u^* to v^* both of length at most $\ell + 1$ such that P^* (resp. Q^*) contains the interior of a principal path P (resp. Q), and $P \neq Q$.

Proof. Let u and v be such that $u^* \in K_u$ and $v^* \in K_v$. Let $P = u \dots v$ and $Q = u \dots v$ be as in the conclusion of Lemma 4.12. By Lemma 6.3, $\{u^*, v^*\} \cup V(P) \setminus \{u, v\}$ and $\{u^*, v^*\} \cup V(Q) \setminus \{u, v\}$ are the desired paths. \Box **Lemma 6.11** If some vertex x of G is adjacent to the ends of a path P of length at most $\ell + 1$ of $G \setminus x$, then x is complete to V(P).

Proof. Otherwise, a shortest cycle in $G[V(P) \cup \{x\}]$ has length at least 4 and at most $\ell+3$. Since $\ell \geq 3$ implies $\ell+3 < 2\ell+1$, this is a contradiction.

6.3 Attaching a vertex to F_1

In this subsection, we show that for all vertices x of $G \setminus F_2$, $N_{V(F_1)}(x)$ is a clique (see Lemma 6.15). In Figure 6, several situations where $N_{V(F_1)}(x)$ is not a clique are represented and we explain informally how they lead to a contradiction. The first figure is an odd 3-template F_0 with its vertices w and w', and here $F_1 = F_0$. Then, vertex x_1 can be included in K_{y_1} , a contradiction to the maximality of F_1 (see Lemma 6.12). The vertex x_2 cannot be included in an existing blown up clique, but it can be added to F_0 to yield a bigger template (see Lemma 6.12). The vertex x_3 can be added to K_{u_6} (see Lemma 6.13). The vertex x_4 can be added to K_{u_1} , but at the expense of modifying the template (see Lemma 6.13). The vertex x_5 can be added to K_{i_3} (see Lemma 6.14).

The vertex x_6 is kind of pathological because it cannot be added to any blown-up clique, and does not increase the template. The idea for this one is to observe that $\{x_6\} \cup V(F_0) \setminus \{y_1, u_6^+\}$ induces a template and that y_1 can be incorporated in the set K_{u_6} and $K_{x_6} = K_{u_6^+} \cup \{x_6\}$ (see Lemma 6.15). Note that in this case, we increase the size of the blowup while decreasing the size of the template.

In each case, we prove that adding x yields a preblowup of F_0 , so that the maximality of F_1 is contradicted.

Lemma 6.12 If $x \in G \setminus F_2$ has no neighbor in I^* , then $N_{V(F_1)}(x)$ is a clique.

Proof. Suppose for a contradiction that $N_{V(F_1)}(x)$ is not a clique.

(1) We may assume that $N_{V(F_1)}(x) \subseteq A^* \cup B^*$.

If x has neighbors in both $A^* \cup B^*$ and $A'^* \cup B'^*$, then consider a path P as in Lemma 6.9 from a neighbor of x in $A^* \cup B^*$ to a neighbor of x in $A'^* \cup B'^*$. By Lemma 6.11, x is complete to V(P). This is a contradiction since x has no neighbor in I^* . Hence x does not have neighbors in both $A^* \cup B^*$ and $A'^* \cup B'^*$, and our claim follows up to symmetry. This proves (1).



Figure 6: Vertices attaching to an odd 3-template

(2) There exist non-adjacent $a, b \in A$ such that x has neighbors in both K_a and K_b .

By Lemma 6.6, since $N_{V(F_1)}(x)$ is not a clique, there should exist two non-adjacent vertices $a, b \in V(F_0)$ such that x has a neighbor $a^* \in K_a$ and a neighbor $b^* \in K_b$. By (1), $a, b \in A \cup B$.

If $a, b \in A$, then our conclusion holds, so we may assume that $b \in B$.

If $a \in A$, then since $ab \notin E(G)$, H_b is anticomplete to a. Let P_a^* be the path induced by $\{a^*\} \cup (V(P_a) \setminus \{a\})$. Let $v \in H_b$. We may assume that $xv \notin E(G)$ for otherwise our claim holds (with a and v). Note that since $ab, av \notin E(G)$, by (c) of blowup, $a^*b^*, a^*v \notin E(G)$. Now, the paths P_a^*, P_v , a^*xb^*v form a hole of length $2\ell + 2$, a contradiction. Hence, we may assume $a \in B$.

Since $ab \notin E(G)$, by Lemma 4.7, $\{a\} \cup H_a$ is anticomplete to $\{b\} \cup H_b$. We may assume that x is anticomplete to $H_a \cup H_b$ for otherwise we may apply the proofs above. Hence, for $u \in H_a$ and $v \in H_b$, the two paths P_u and P_v together with the path ua^*xb^*v form a hole of length $2\ell + 3$. This proves (2).

Now the sets K_u for all $u \in A \cup A' \cup I$, $B^* \cup \{x\}$ and B'^* form a preblowup of F_0 . All conditions are easily checked. In particular x satisfies condition (b) by (1) and (b2) by (2)). So, by Lemma 5.6, $G[V(F_1) \cup \{x\}]$ is a proper blowup of some ℓ -template with k principal paths. This contradicts the maximality of F_1 .

Lemma 6.13 If there exist $x \in V(G) \setminus V(F_2)$ and $u \in A$ such that x has neighbors in both K_u and K_{u^+} and is anticomplete to $K_{u^{++}}$, then $N_{V(F_1)}(x)$ is a clique.

Proof. Suppose for a contradiction that $N_{V(F_1)}(x)$ is not a clique.

(1) x is anticomplete to $A'^* \cup B'^* \cup (I^* \setminus K_{u^+})$.

If x has a neighbor t^* in some K_t such that $t \in (A' \cup I) \setminus \{u^+\}$, then note that $t \neq u^{++}$ by assumption. Let C be a principal hole that contains t and u. There is a contradiction to Lemma 6.8 because by (c) of blowup u, u^+ and t cannot be consecutive along C.

It remains to prove that x is anticomplete to B'^* . Otherwise, x has a neighbor $t \in B'^*$. Consider a path P from t to the neighbor of x in K_u as in Lemma 6.9 and let Q be the principal path whose interior is contained in P. By Lemma 6.11, x is complete to V(P). This is a contradiction because if $Q = P_u$ then x is anticomplete to $K_{u^{++}}$, and if $Q \neq P_u$ then we already proved that x is anticomplete to $(A'^* \cup I^*) \setminus K_{u^+}$. This proves (1).

From here on, u^* and u^{+*} are neighbors of x in respectively K_u and K_{u^+} . Note that x has a neighbor $y^* \in K_y$ for some $y \in A \cup B \setminus \{u\}$, for otherwise, by (1), $N_{V(F_1)}(x) \subseteq K_u \cup K_{u^+}$ and by Lemma 6.6, $N_{V(F_1)}(x)$ is a clique, a contradiction.

(2) If $w \in B$, then x has a neighbor $w^* \in B^*$ that is complete to A^* .

We may assume that x is non-adjacent to w, for otherwise by condition (h) of blowups, we may choose $w^* = w$. In particular $y^* \neq w$.

We claim that we may assume that y^* has a non-neighbor v^* such that $v^* \in K_v, v \in A$ and $v \neq u$.

If $y^* \in B^*$, this is because we may assume that y^* has a non-neighbor $v^* \in A^*$ (so $v^* \in K_v$ for some $v \in A$) for otherwise we choose $w^* = y^*$ from the start. It remains to check that $u \neq v$. This is because if u = v, then there exists a path Q of length 1, 2 or 3 from x to v^* with interior in K_{u^+} (through xv^* , u^+ , u^{+*} or u^+u^{+*}). Hence, xQv^*wy^*x is a hole of length 4, 5 or 6, a contradiction.

If $y^* \in A^*$, then $u^*y^* \in E(G)$ for otherwise, $\{x, y^*, w, u^*\}$ induces a C_4 . By condition (c) of blowups, $uy \in E(G)$. It follows that none of u and y is isolated in G[A], so the existence of v^* follows from Lemma 4.17 that guarantees the existence of isolated vertices in G[A] since $w \in B$ by assumption.

So, our claim is proved. Note that $xv^* \notin E(G)$ for otherwise $\{x, y^*, w, v^*\}$ induces a C_4 . Now either $xy^*wv^*v^+P_vv'w'u'P_uu^{++}u^{+*}x$ is a hole of length $2\ell + 3$ (in case $u'v' \notin E(G)$) or $xy^*wv^*v^+P_vv'u'P_uu^{++}u^{+*}x$ is a hole of length $2\ell + 2$ (in case $u'v' \in E(G)$). In both cases we get a contradiction. This proves (2).

(3) $N_A(x) \setminus \{u\} = N_A(u).$

If there exists $v \in N_A(x) \setminus N_A[u]$, then $v P_v v' u' P_u u^{++} u^{+*} x v$ is a hole of length 2ℓ , a contradiction.

Conversely, suppose there exists $v \in N_A(u) \setminus N_A(x)$. We claim that there exists a path Q of length 2 from x to some $z \in N_A(u)$ with interior in $(A^* \cup B^*) \setminus (K_u \cup K_z)$.

If $w \in B$, then we may choose z = v and $Q = xw^*z$ by (2).

Otherwise, $w \in A$. So, by Lemma 4.17, G[A] contains at least two universal vertices. So, let $t \in A \setminus \{u, v\}$ be adjacent to u and v (if u and v are the universal vertices of G[A], t can be any vertex of $A \setminus \{u, v\}$ and otherwise choose t to be a universal vertex).

If x has a neighbor t^* in K_t , then we choose $Q = xt^*v$. So, suppose x is anticomplete to K_t (in particular, $y \neq t$). If x has a neighbor v^* in K_v , then we choose $Q = xv^*t$. So, suppose x is anticomplete to K_v (in particular, $y \neq v$). Now, by the way we chose v and t, one of v or t is a universal vertex of G[A] and therefore a universal vertex of $G[A^* \cup B^*]$. So, we may choose $Q = xy^*v$ or $Q = xy^*t$.

So, our claim is proved. Hence $z'P_z zQxu^{+*}u^{++}P_uu'w'z'$ is a hole of length $2\ell + 2$, a contradiction. This proves (3).

(4) x is complete to K_u .

Suppose there exists $r \in K_u$ such that $rx \notin E(G)$. We claim that x and r have a common neighbor z in $(A^* \cup B^*) \setminus K_u$.

If $w \in B$, then $rw^* \in E(G)$ by (2) so we may choose $z = w^*$. If $w \in A$, then by Lemma 4.17, some vertex $z \in A \setminus \{u\}$ is a universal vertex of G[A], and by (3), z is adjacent to x. So, z exists as claimed.

If $xu^+ \in E(G)$ then $\{r, z, u^+, x\}$ induces a C_4 , a contradiction. Hence $xu^+ \notin E(G)$. Now by condition (e) of blowups, either $\{x, z, r, u^{+*}\}$ induces a C_4 or $\{x, z, r, u^+, u^{+*}\}$ induces a C_5 . This proves (4).

Now, the sets K_v for all $v \in (A \setminus u) \cup I \cup A'$, $K_u \cup \{x\}$, B^* and B'^* form a preblowup of F_0 . All conditions are easy to check. In particular, $K_u \cup \{x\}$ is a clique by (4), conditions (a), (b) and (i) follows from (1), condition (a1) from (3), condition (b1) from (2) and condition (a2) from our assumptions.

Hence, by Lemma 5.6 $G[V(F_1) \cup \{x\}]$ is a proper blowup of some twinless odd ℓ -template with k principal paths that is an induced subgraph of G a contradiction to the maximality of F_1 .

Lemma 6.14 If $x \in V(G) \setminus V(F_2)$ has no neighbor in $B^* \cup B'^*$, then $N_{V(F_1)}(x)$ is a clique.

Proof. Suppose for a contradiction that $N_{V(F_1)}(x)$ is not a clique. By Lemma 6.12, x has neighbors in I^* . So x has a neighbor in a clique blown up from an internal vertex of some principal path $P_v = v \dots v'$. Let a (resp. b) be the vertex of P_v closest to v (resp. to v') along P_v and such that x has a neighbor in K_a (resp. K_b).

Suppose first that a = b (so $a \in I$). Then x has a neighbor in some K_y with $y \in V(F_0) \setminus \{a\}$, and since by assumption x has no neighbor in $B^* \cup B'^*$, $y \in A \cup A' \cup I$. So, y and a are non-adjacent members of some

principal hole. By Lemma 6.8, x has a neighbor in some clique K_d where d is adjacent to a = b, a contradiction to a = b.

Suppose now that $ab \in E(G)$. If both a and b are internal vertices of P_v , then as in the previous paragraph, we may deduce from Lemma 6.8 that $N_{V(F_1)}(x) \subseteq K_a \cup K_b$. So, by Lemma 6.6, $N_{V(F_1)}(x)$ is a clique, a contradiction. It follows that at least one of a or b is an end of P_v . Up to symmetry, we may assume that a = v and $b = v^+$. Note that x is then anticomplete to $K_{v^{++}}$. Hence, by Lemma 6.13, $N_{V(F_1)}(x)$ is a clique, a contradiction.

Hence, $a \neq b$ and $ab \notin E(G)$. So, by Lemma 6.8, a and b have a common neighbor u in P_v . So, a, u and b are consecutive along P_v (in particular, $u \in I$).

(1) x is complete to K_u .

Otherwise, let $u^* \in K_u$ be a non-adjacent to x. There exists a path Q_a of length 2 or 3 from u^* to x with interior in K_a (either xa^*u^* , or xa^*au^* for some a^* in K_a). There exists a similar path Q_b . So, Q_a and Q_b form a hole of length 4, 5 or 6, a contradiction. This proves (1).

(2) x is anticomplete to $V(F_1) \setminus (K_a \cup K_u \cup K_b)$.

This follows from Lemma 6.8 and from the fact that x is anticomplete to $B^* \cup B'^*$. This proves (2).

(3) x has neighbors in each of K_a , K_b .

This follows from the definition of a and b. This proves (3).

Now the sets K_v for all $v \in (A \cup A' \cup I) \setminus \{u\}$, $K_u \cup \{x\}$, B^* and B'^* form a preblowup of F_0 . All conditions are easily checked, in particular $K_u \cup \{x\}$ is a clique by (1), it satisfies condition (i) by (2) and condition (i1) by (3).

Hence by Lemma 5.6, $G[V(F_1) \cup \{x\}]$ is a proper blowup of some twinless odd ℓ -template with k principal paths that is an induced subgraph of G. This contradicts the maximality of F_1 .

Lemma 6.15 For all vertices x of $G \setminus F_2$, $N_{V(F_1)}(x)$ is a clique.

Proof. Suppose for a contradiction that $N_{V(F_1)}(x)$ is not a clique.

(1) There exists a principal path $P_u = u \dots u'$ of F_0 such that x is anticomplete to $I^* \setminus \bigcup_{v \in V(P_u)} K_v$.

Otherwise, there exist two principal paths P and Q of F_0 , a in the interior of P and b in the interior of Q such that x has neighbors in both K_a and K_b .

Note that P and Q are in some principal hole C of F_0 . By Lemma 6.8, a and b have a common neighbor c in C. This contradicts a and b being in the interior of distinct principal paths. This proves (1).

(2) We may assume that x has no neighbor in $B^{\prime*}$ and has a neighbor $y^* \in K_y$ where $y \in B$.

Suppose that x has a neighbor $u^* \in B^*$ and a neighbor $v^* \in B'^*$. Let P and Q be like in Lemma 6.10. By Lemma 6.11, x is complete to both V(P) and V(Q). In particular, x has neighbors in the interior of two distinct principal paths, a contradiction to (1). So, up to symmetry, we may assume that x has no neighbor in B'^* . Hence, by Lemma 6.14, x has neighbors in B^* . This proves (2).

(3) x is adjacent to u, u^+ and has a neighbor in $K_{u^{++}}$. Moreover, x is anticomplete to $(A^* \cup I^* \cup A'^* \cup B'^*) \setminus (K_u \cup K_{u^+} \cup K_{u^{++}})$.

By Lemma 6.12, x has at least one neighbor in I^* and by (1), such a neighbor is in a clique blown up from an internal vertex of P_u . So, let v be the vertex of P_u closest to u' along P_u such that x has a neighbor $v^* \in K_v$. So $v \neq u$ and $v \in A' \cup I$. We set $Q = y^* u P_u v$ if $y^* u \in E(G)$ and $Q = y^* w u P_u v$ otherwise. Let Q^* be the path induced by $\{v^*\} \cup (V(Q) \setminus \{v\})$ and observe that Q^* has length at most $\ell + 1$. By Lemma 6.11, x is complete to Q^* . If $v \notin \{u^+, u^{++}\}$, then x has neighbors in at least 4 cliques blown up from vertices of P_u and this contradicts Lemma 6.8. If $v = u^+$, x is adjacent to u (since x is complete to Q^*) and anticomplete to $K_{u^{++}}$, so by Lemma 6.13, $N_{V(F_1)}(x)$ is a clique, a contradiction. So, $v = u^{++}$, meaning that x is adjacent to u and u^+ , and is anticomplete to $I^* \setminus (K_{u^+} \cup K_{u^{++}})$ by (1).

If x has neighbors in some K_a for $a \in A \setminus \{u\}$ then x and $C_{u,a}$ contradict Lemma 6.8. Hence x is anticomplete to $A^* \setminus \{K_u\}$.

By (2), x is anticomplete to B'^* . It remains to check that x is anticomplete to $A'^* \setminus K_{u^{++}}$. So, suppose x has a neighbor z^* in some K_z where $z \in A' \setminus \{u^{++}\}$. Then a principal hole that contains z and u contradicts Lemma 6.8. This proves (3).

Let u^{++*} be a neighbor of x in $K_{u^{++}}$ and P_u^* be the path induced by $(V(P_u) \setminus \{u^{++}\}) \cup \{u^{++*}\}.$

(4) For every $z \in B$ such that x is adjacent to some z^* in K_z we have $N_A(z) = N_A[u]$ (in particular $N_A(y) = N_A[u]$).

Suppose there exists $v \in N_A(z) \setminus N_A[u]$. By condition (d) or (f) of blowups, $vz^* \in E(G)$. So, by (3), $xz^*vP_vv'u'P_u^*u^{++*}x$ is a hole of length 2ℓ , a con-

tradiction. This proves that $N_A(z) \subseteq N_A[u]$. In particular, u has at least one neighbor in H_z , so by condition (e5) of templates, $uz \in E(G)$.

Suppose there exists $v \in N_A(u) \setminus N_A(z)$ (so z and v are not universal vertices of $G[A \cup B]$). By condition (c) of blowups, $vz^* \notin E(G)$. By (3), $xv \notin E(G)$. Hence $xd \in E(G)$ for every universal vertex d of $G[A \cup B]$, for otherwise $xz^*dvP_vv'w'u'P_u^*u^{++*}x$ is a hole of length $2\ell + 2$.

Now, by (3) and Lemma 4.17, $w \in B$. So, there exists an isolated vertex $c \in A$. Again by (3), $xc \notin E(G)$ and $xwcP_cc'u'P_u^*u^{++*}x$ is a hole of length 2ℓ , a contradiction. This proves (4).

$$(5) N_{F_1}(x) \subseteq K_{u^{++}} \cup K_{u^+} \cup K_u \cup K_y$$

By (3) $N_{F_1}(x) \subseteq K_{u^{++}} \cup K_{u^+} \cup K_u \cup B^*$. Suppose there exists $z^* \in K_z$ such that $xz^* \in E(G)$ and $z \in B \setminus \{y\}$. By (4), $N_A(z) = N_A[u]$ and $N_A(y) = N_A[u]$. So, by Lemma 4.15, y and z are twins of F_0 , a contradiction. This proves (5).

(6) $y \neq w$.

If y = w, then $w \in B$. So by Lemma 4.17, there exist isolated vertices in G[A]. But by (4), $N_A(w) = N_A[u]$ so u is a universal vertex of G[A], so G[A] has a universal vertex and an isolated vertex, a contradiction. This proves (6).

(7) $N_{K_u}(x)$ is complete to $N_A[u]$.

By (4), $N_A(y) = N_A[u]$. The result follows from conditions (d) and (f) of blowups. This proves (7).

(8) x is complete to K_{u^+} .

By (3), $ux \in E(G)$. Suppose for a contradiction that there exists $u^{+*} \in K_{u^+}$ non-adjacent to x. By condition (e) of blowups, $u^{+*}u, u^{+*}u^{++} \in E(G)$. Hence $xu^{++} \notin E(G)$ for otherwise $\{x, u^{++}, u^{+*}, u\}$ induces a C_4 . But now, either $\{x, u^{++*}, u^{+*}, u\}$ induces a C_4 (if $u^{+*}u^{++*} \in E(G)$) or $\{x, u^{++*}, u^{+*}, u\}$ induces a C_5 (if $u^{+*}u^{++*} \notin E(G)$), a contradiction. This proves (8).

(9) $K_u \cup K_y$ is a clique.

Since by (4) $N_A(y) = N_A[u]$, *u* cannot be an isolated vertex of H_y . Hence, *uy* is a solid edge. So, by condition (d) of blowups, K_u is complete K_y . This proves (9).

We define $B_0 = B^* \setminus N_{K_y}(x)$.

Now the sets K_v for all $v \in (A \cup I \cup A') \setminus \{u, u^+\}, K_u \cup N_{K_y}(x), K_{u^+} \cup \{x\}, B_0$ and B'^* form a preblowup of F_0 . All conditions are easy to check. In particular, $K_u \cup N_{K_y}(x)$ is a clique by (9), $K_{u^+} \cup \{x\}$ is a clique by (8), conditions (a), (b) and (i) follows from (5), condition (a1) from (7), condition (a2) holds because x is complete to $N_{K_y}(x)$, condition (i1) follows from (3) and condition (b1) holds because (6) implies that if $w \in B$ then $w \in B_0$.

Hence, by Lemma 5.6, $G[V(F_1) \cup \{x\}]$ is a proper blowup of some twinless odd ℓ -template with k principal paths that is an induced subgraph of G, a contradiction to the maximality of F_1 .

6.4 Attaching a component

Lemma 6.16 If D is a connected component of $G \setminus F_2$, then N(D) is a clique.

Proof. Suppose that N(D) is not a clique. By Lemma 6.2, $N_{V(F_1)}(D)$ is not a clique. So, there exist a and b in D such that $N_{V(F_1)}(a) \cup N_{V(F_1)}(b)$ is not a clique, and a path P from a to b in D. We choose a and b subject to the minimality of the length of P. By Lemma 6.15, $a \neq b$ (so P has length at least 1).

We set $S_a^* = N_{V(F_1)}(a)$ and $S_b^* = N_{V(F_1)}(b)$. By Lemma 6.15, S_a^* and S_b^* are both cliques. Note that possibly $S_a^* \cap S_b^* \neq \emptyset$. We denote by $\operatorname{int}(P)$ the set of the internal vertices of P. We set $S_{\circ}^* = N_{V(F_1)}(\operatorname{int}(P))$.

We set $S_a = \{t \in V(F_0) : S_a^* \cap K_t \neq \emptyset\}$. We define S_b and S_o similarly. Note that S_a is possibly not included in S_a^* , and the same remark holds for S_b and S_o .

(1) There exist non-adjacent $x_a^* \in S_a^*$ and $x_b^* \in S_b^*$. Moreover, for all such x_a^* and x_b^* , $x_a^* aPbx_b^*$ is a path.

The existence of x_a^* and x_b^* follows from the definition of a and b, and $x_a^* a P b x_b^*$ is a path because of the minimality of P. This proves (1).

(2) $S_a^* \cup S_o^*$ and $S_b^* \cup S_o^*$ are cliques (in particular, S_o^* is a (possibly empty) clique of F_1 that is complete to both $S_a^* \setminus S_o^*$ and $S_b^* \setminus S_o^*$).

If $S_a^* \cup S_o^*$ is not a clique, then let x^*y^* be a non-edge in $S_a^* \cup S_o^*$. Since S_a^* is a clique by Lemma 6.15, we may assume $y^* \in S_o^*$. By definition of S_o^* , y^* has a neighbor in int(P), and then x^*, y^* and some subpath of P contradict the minimality of P.

The proof is similar for $S_b^* \cup S_o^*$. This proves (2).

Note that while $S_a^* \cup S_b^*$ is not a clique by assumption, it might be that $S_a \cup S_b$ is a clique (for instance when $S_a = \{u\}$, $S_b = \{v\}$ and uv is an optional edge of F_0).

(3) $S_a \cup S_\circ$ and $S_b \cup S_\circ$ are cliques of F_0 (in particular, S_a and S_b are (non-empty) cliques of F_0 and S_\circ is a (possibly empty) clique of F_0 that is complete to both $S_a \setminus S_\circ$ and $S_b \setminus S_\circ$).

If $S_a \cup S_\circ$ is not a clique, then let xy be a non-edge of $S_a \cup S_\circ$. Since $x \in S_a \cup S_\circ$, there exists $x^* \in K_x \cap (S_a^* \cup S_\circ^*)$ and $y^* \in K_y \cap (S_a^* \cup S_\circ^*)$. By condition (c) of blowups, since $xy \notin E(G)$, K_x is anticomplete to K_y . So, $x^*y^* \notin E(G)$, a contradiction to (2).

The proof is similar for $S_b \cup S_{\circ}$. This proves (3).

(4) If a hole C of F_1 contains two non adjacent vertices $x \in S_a^*$ and $y \in S_b^*$, then P and C form a pyramid $\prod_{C,x,y}$. More specifically, C contains a vertex z such that either:

- $S_a^* \cap V(C) = \{x, z\}, S_b^* \cap V(C) = \{y\}$; the apex of $\Pi_{C,x,y}$ is y, its triangle is axz, and its three paths, all of length ℓ , are the path from x to y in $C \setminus z$, the path from y to z in $C \setminus x$, and the path from a to y obtained by adding the edge by to P; or
- $S_b^* \cap V(C) = \{y, z\}, S_a^* \cap V(C) = \{x\}$; the apex of $\Pi_{C,x,y}$ is x, its triangle is by z, and its three paths, all of length ℓ , are the path between y and x in $C \setminus z$, the path from z to x in $C \setminus y$, and the path from b to x obtained by adding the edge ax to P.

Note that since S_a^* is a clique, $S_a^* \cap V(C)$ contains x and at most one other vertex which should be adjacent to x. The same holds for S_b^* and y.

Let us assume that $S_{\circ}^{*} \cap V(C) \neq \emptyset$. Then by (2), there exists a unique vertex $t \in S_{\circ}^{*} \cap V(C)$, $S_{a}^{*} \cap V(C) \subseteq \{x,t\}$ and $S_{b}^{*} \cap V(C) \subseteq \{y,t\}$. Hence C and P form a proper wheel centered at t, a contradiction to Lemma 3.10. So, $S_{\circ}^{*} \cap V(C) = \emptyset$.

If a and b have a common neighbor t in C, then x and y are the two neighbors of t in C and so, C and P form a proper wheel centered at t, again a contradiction to Lemma 3.10. So the neighborhoods of a and b in C are disjoint.

From this, we obtain that C and P form a theta, a prism or a pyramid. So, by Lemma 3.10, C and P form a pyramid whose three paths have length ℓ . This can happen only if we are in one of the two cases described in (4). This proves (4). (5) $S_a \cap I = S_b \cap I = \emptyset$.

Otherwise, up to symmetry, $S_a \cap I \neq \emptyset$. So, there exists a principal path $P_u = u \dots u'$ of F_0 whose interior intersects S_a . By (3), S_a is a clique, so $1 \leq |S_a| \leq 2$ and $S_a \subseteq V(P_u)$. We now break into three cases.

Case 1: $S_b \subseteq V(P_u)$.

By (1) there exist vertices x_a and x_b of P_u such that there exist non adjacent vertices $x_a^* \in S_a^* \cap K_{x_a}$ and $x_b^* \in S_b^* \cap K_{x_b}$.

We first show that there exist such x_a and x_b that are not adjacent. Otherwise, and since $S_a, S_b \subseteq V(P_u)$, we have that $S_a \cup S_b = \{x_a, x_b\}$. By replacing x_a and x_b by x_a^* and x_b^* in any principal hole C containing P_u we obtain a path P_C of length 2ℓ and $V(P_C) \cup V(P)$ induces a hole of length at least $2\ell + 3$, a contradiction. So we may assume that x_a and x_b are not adjacent.

Let C be any principal hole of F_0 that contains P_u . By Lemma 6.3, $\{x_a^*, x_b^*\} \cup (V(C) \setminus \{x_a, x_b\})$ induces a hole C^* . Let us apply (4) to C^* , x_a^* and x_b^* . We obtain that the shortest path in C^* between x_a^* and x_b^* has length ℓ . However x_a^* and x_b^* both belong to the path of length $\ell - 1$, contained in C^* , which is obtained from P_u by replacing x_a by x_a^* and x_b by x_b^* , a contradiction.

Case 2: S_b contains a vertex of some principal path P_v distinct from P_u . Up to symmetry, since S_b is a clique (by (3)), we assume that b is anticomplete to $K_{v'}$.

Let y be the vertex of P_u closest to u' such that a has a neighbor $y^* \in K_y$. Let z be the vertex of P_v closest to v such that b has a neighbor $z^* \in K_z$. Possibly y = u' and z = v, but $y \neq u$ since a has a neighbor in I^* by assumption, and $z \neq v'$ since b is anticomplete to $K_{v'}$. In particular, $yz \notin E(G)$ and by condition (c) of blowups, $y^*z^* \notin E(G)$.

Let C be the principal hole of F_0 that contains P_u and P_v . By Lemma 6.3, $\{y^*, z^*\} \cup (V(C) \setminus \{y, z\})$ induces a hole C^* . Applying (4) to C^* , y^* and z^* , we obtain that P has length $\ell - 1$ and that y^* and z^* are at distance ℓ on C^* . Hence y^* and z^* have no common neighbor in F_0 and $S_0 = \emptyset$ by (3). We denote by P_u^* the path obtained from P_u by replacing y by y^* and by P_v^* the path obtained from P_v by replacing z by z^* . Let P^* be the path $vP_v^*z^*bPay^*P_u^*u'$ (in case z = v one should replace $vP_v^*z^*$ by z^* , and in case y = u' one should replace $y^*P_u^*u'$ by y^*). The length of P^* is at least $\ell + 1$.

Consider now any principal path P_r for $r \in A \setminus \{u, v\}$. Depending on the adjacencies of r with u and v, one of $rvP^*u'r'P_rr$ or $rwvP^*u'r'P_rr$ or $rvP^*u'w'r'P_rr$ or $rwvP^*u'w'r'P_rr$ (with possibly u' replaced by y^* when u' = y) is a cycle of length at least $2\ell + 2$ with at most one chord that must be br (observe that ar' cannot be an edge since $S_a \subseteq V(P_u)$). The only possibility which avoids a hole of forbidden length is if z = v, y = u' and br, u'r', vr are edges of G. This proves that v is complete to $A \setminus \{u, v\}$ and u' is complete to $A' \setminus \{u', v'\}$.

Hence, G[A] has at most one isolated vertex (namely u), and G[A'] has at most one isolated vertex (namely v'). This contradicts (A, B, A', B', I, w, w') being a proper ℓ -partition of F_0 .

Case 3: we are neither in Case 1 nor in Case 2.

Since we are not in Case 1, S_b contains a vertex of $F_0 \setminus P_u$, and since we are not in Case 2, this vertex must be in $B \cup B'$. Up to symmetry, we assume that $S_b \cap B \neq \emptyset$. Since S_b is a clique (by (3)), $S_b \cap (B' \cup A' \cup I) = \emptyset$. Since we are not in Case 2, $S_b \cap (A \setminus \{u\}) = \emptyset$. Hence, $S_b \subseteq B \cup \{u\}$ and there exists $x \in B \cap S_b$. Let $x^* \in K_x \cap S_b^*$.

Let u_a be the vertex of S_a which is the closest to u in P_u and let u'_a be the vertex of S_a which is the closest to u' in P_u . Notice that, since S_a is a clique (by (3)), either $u_a = u'_a$ or $u_a u'_a$ is an edge. So it may be that $u_a = u$ or $u'_a = u'$ but since $S_a \cap I \neq \emptyset$ we know that $u_a \neq u'$ and $u'_a \neq u$. Let now $u^*_a \in K_{u_a} \cap S^*_a$ and $u'^*_a \in K_{u'_a} \cap S^*_a$. We denote by P^*_u the path obtained from P_u by replacing u_a by u^*_a and, in case $u_a \neq u'_a$, by replacing u'_a by u'^*_a . Notice that if $u^*_a \neq u'^*_a$ then $u^*_a u'^*_a \in E(G)$ since S^*_a is a clique.

Suppose that $u'_a = u^+$, where u^+ is the neighbor of u in P_u . Since H_x contains at least two vertices there exists $v \in H_x \setminus \{u\}$. By (3) and the fact that P contains at least one edge, depending on the adjacency of u and v, one of $aPbx^*vP_vv'u'P_u^*u'_a^*a$ or $aPbx^*vP_vv'w'u'P_u^*u'_a^*a$ is a hole of length at least $2\ell+2$, a contradiction. Hence from now on, we may assume that $u_a \neq u$ (hence a is not adjacent to u) and that if $u_a = u^+$ then $u'_a \neq u_a$. Now by (3) we get that $S_o = \emptyset$.

Suppose that x is adjacent to u in F_0 . Depending on whether b is adjacent to u not, one of $u_a^* a P b u P_u^* u_a^*$ or $u_a^* a P b x^* u P_u^* u_a^*$ is a hole, implying that P has length at least ℓ . Let us choose any vertex $v \in H_x$ distinct from u (since H_x has cardinality at least 2, such a vertex do exist). Then $a P b x^* v P_v v'(w') u' P_u u_a'^* a$ is a hole of length at least $2\ell + 2$, a contradiction. Hence, from here on, we may assume that no vertex in $B \cap S_b$ is adjacent to u.

So x is not adjacent to u in F_0 . Hence $x^* \neq w$, $u \neq w$ and $w \notin S_b$. Then to avoid a C_4 bx^*wub , b is not adjacent to u and $u_a^*aPbx^*wuP_u^*u_a^*$ is a hole implying that P has length at least $\ell - 1$. So, for any $v \in H_x$, the hole $x^*bPau'_a^*P_u^*u'v'P_vvx^*$ (in case $u'_a = u'$ one should replace $u'_a^*P^*u'$ by u'_a^*) has length at least $2\ell + 2$, a contradiction. This proves (5).

(6) We may assume that $S_a \subseteq A \cup B$ and $S_b \subseteq A' \cup B'$.

Otherwise, by (5) and since S_a and S_b are cliques (by (3)), we may assume that $S_a, S_b \subseteq A \cup B$.

We claim that there exist non-adjacent vertices $x^* \in S_a^*$ and $y^* \in S_b^*$, and a path Q^* from x^* to y^* of length at least $2\ell - 1$ that forms a hole together with P. This is a contradiction because it implies that P has length at most 0. So, to conclude the proof, it remains to prove the existence of Q^* .

By (1), there exist non-adjacent $x_a^* \in S_a^*$ and $x_b^* \in S_b^*$. Let x_a and x_b be the vertices of F_0 such that $x_a^* \in K_{x_a}$ and $x_b^* \in K_{x_b}$. Note that possibly $x_a x_b$ is an edge, but this happens only if $x_a x_b$ is an optional edge of F_0 (since $x_a^* x_b^*$ is not an edge). We break into three cases.

Case 1: $x_a, x_b \in A$.

Then $x_a x_b \notin E(G)$ (otherwise it would be a solid edge of F_0), so from the definition of templates, there exists a path Q of length $2\ell - 1$ from x_a to x_b whose interior is in $I \cup A'$. By Lemma 6.3, $\{x_a^*, x_b^*\} \cup (V(Q) \setminus \{x_a, x_b\})$ induces the path Q^* that we are looking for. Note that Q^* and P form a hole by (2), our assumption that $S_a, S_b \subseteq A \cup B$, and (3).

Case 2: $x_a \in A$ and $x_b \in B$.

Whether $x_a x_b$ is an optional edge or a non-edge, an immediate consequence of the definition of a template is that there exists a vertex $z \in H_{x_b}$ that is non-adjacent to x_a . We may furthermore assume that $z \notin S_b$ since else we are in the same situation as in Case 1. By definition of a template, there exists a path Q_0 of length $2\ell - 1$ between x and z whose interior is in $I \cup A'$. Then $x_b z Q_0 x_a$ is a path of length 2ℓ and by replacing in this path x_a and x_b by respectively x_a^* and x_b^* , we obtain by Lemma 6.3 a path Q^* of the same length. Note that Q^* and P form a hole by (2), (3), our assumption that $S_a, S_b \subseteq A \cup B, z \notin S_b$ and $z \notin S_a$ (since S_a is a clique).

Case 3: $x_a, x_b \in B$.

Then $x_a x_b \notin E(G)$ (otherwise it would be a solid edge of F_0). Hence, by Lemma 4.7, $H_{x_a} \cup \{x_a\}$ is anticomplete to $H_{x_b} \cup \{x_b\}$. So, let $u_a \in H_{x_a}$ and $u_b \in H_{x_b}$, there exists then a path $Q_0 = u_a \dots u_b$ of length $2\ell - 1$ with interior in $I \cup A'$. By Lemma 6.3, $Q^* = x_a^* u_a Q_0 u_b x_b^*$ is also a path, it is of length $2\ell + 1$. We may assume that $u_a \notin S_a$ and $u_b \notin S_b$ since else we are in the same situation as in Case 2. Now, by (2) and (3), Q^* and P form a hole of length at least $2\ell + 4$.

This proves (6).

(7) $S_{\circ} = \emptyset$.

By (6) and (3), if $S_{\circ} \neq \emptyset$, then $\ell = 3$, and there exists a unique principal path $P_u = u \dots u'$ of F_0 such that $S_a = \{u\}$, $S_b = \{u'\}$ and $S_{\circ} = \{c\}$ where c is the unique internal vertex of P_u . Let $u^* \in K_u \cap S_a^*$, $c^* \in S_{\circ}^*$ and $u'^* \in K_{u'} \cap S_b^*$. Observe that by (2) $c^*u^*, c^*u'^* \in E(G)$.

Let $P_v = v \dots v'$ be a principal path distinct from P_u and suppose up to symmetry that $uv \in E(G)$. Now by (3), P_v , P, u^* , u'^* , w' and c^* form a proper wheel centered at c^* , a contradiction to Lemma 3.10. This proves (7).

(8) P has length $\ell - 1$, or P has length $\ell - 2$ and we may assume that $S_a \cap A = \emptyset$.

By (1) and (6), consider non-adjacent $x^* \in S_a^*$ and $y^* \in S_b^*$ where $x^* \in K_x \cap S_a^*$ and $y^* \in K_y \cap S_b^*$ for $x \in S_a \cap (A \cup B)$ and $y \in S_b \cap (A' \cup B')$.

If $x \in A$ and $y \in A'$, then let C be a principal hole that contains x and y. By Lemma 6.3, $\{x^*, y^*\} \cup (V(P) \setminus \{x, y\})$ induces a hole C^* . We may apply (4) to C^* , x^* and y^* . It follows that P has length $\ell - 1$. By symmetry we may therefore assume from here on that $S_a \cap A = \emptyset$.

Let y'^* be a vertex in S_b^* which is the closest to x^* in F_1 . By Lemma 6.9, there exists a path Q in F_1 from x^* to y'^* of length ℓ or $\ell + 1$. From our assumption on y'^* we get that Q and P form a hole (since $S_o = \emptyset$ by (7)). Therefore, if Q has length ℓ , then P has length $\ell - 1$ and if Q has length $\ell + 1$, then P has length $\ell - 2$. This proves (8).

We may now conclude the proof.

If P has length $\ell - 1$, then we set $A_0 = A \cup \{a\}$, $A'_0 = A' \cup \{b\}$ and $I_0 = I \cup \operatorname{int}(P)$. We claim that (A_0, B, A'_0, B', I_0) is an ℓ -pretemplate partition of $G[A_0 \cup B \cup A'_0 \cup B' \cup I_0]$. All conditions are easily checked to hold (in particular conditions (a), (b) and (g) are satisfied because by (6), a (resp. b) has a neighbor in $G[A \cup B]$ (resp. $G[A' \cup B']$), condition (e) holds by (7) and conditions (h) and (i) hold because they hold in F_0). Then, by Lemma 4.14, G contains an odd ℓ -template with k + 1 principal paths, a contradiction to the maximality of k.

So, by (8), P has length $\ell - 2$ and we may assume that $S_a \cap A = \emptyset$ and $S_a \cap B \neq \emptyset$ (recall that by (6), $S_a \subseteq A \cup B$ and $S_b \subseteq A' \cup B'$). Let us choose $x \in S_a \cap B$ such that H_x is maximal (note that x is unique because $S_a \cap B$ is a clique and F_0 is twinless). Let $x^* \in K_x \cap S_a^*$. We set $A_0 = A \cup \{x^*\}$, $B_0 = B \setminus S_a$, $A'_0 = A' \cup \{b\}$ and $I_0 = I \cup \operatorname{int}(P) \cup \{a\}$. Note that the path x^*aPb has length $\ell - 1$ and has interior in I_0 . We break into two cases.

Case 1: *b* has a neighbor in $A' \cup B'^*$.

We claim that in that case $(A_0, B_0, A'_0, B'^*, I_0)$ is an ℓ -pretemplate partition of $G[A_0 \cup B_0 \cup A'_0 \cup B'^* \cup I_0]$. All conditions are easily checked to hold (in particular condition (g) is satisfied for $A_0 \cup B_0$ because if x = w, then x^* is complete to $(A_0 \cup B_0) \setminus \{x^*\}$, and otherwise, by the maximality of H_x , $w \in A_0 \cup B_0$, condition (g) is satisfied for $A'_0 \cup B'^*$ because b has a neighbor in $A' \cup B'^*$ and by the rules of the blowup, conditions (a), (b), (h) and (i) hold because they hold in F_0 and by the rules of the blowup). Then, by Lemma 4.14, G contains an odd ℓ -template in with k + 1 principal paths, a contradiction to the maximality of k.

Case 2: b has no neighbor in $A' \cup B'^*$.

Then, by (6) there exists $x'^* \in K_{x'} \cap S_b^*$ for some $x' \in A'$.

Let $A'_1 = (A'_0 \cup \{x'^*\}) \setminus \{x'\}$. If $w' \in B'$ we set $B'_1 = \{w'\}$ and else we set $B'_1 = \emptyset$. We claim that $(A_0, B_0, A'_1, B'_1, I_0)$ is an ℓ -pretemplate partition of $G[A_0 \cup B_0 \cup A'_1 \cup B'_1 \cup I_0]$.

Most conditions are easily checked to hold as in the previous case. Notice that conditions (g) and (i) hold because $x^{\prime*}$ is by definition adjacent to b and by the rules of the blowup, $G[A'_1 \setminus \{b\}]$ is isomorphic to $G[A'_0 \setminus \{b\}]$ and $x^{\prime*}$ is adjacent to w'. Then, by Lemma 4.14, G contains an odd ℓ -template in with k + 1 principal paths, a contradiction to the maximality of k. \Box

6.5 End of the proof

We may now conclude the proof of Lemma 6.1. If $G \setminus F_1$ is empty, then conclusion (a) holds. If $G \setminus F_1$ is non-empty and $G \setminus F_2$ is empty, then conclusion (b) holds. Otherwise, we consider a connected component D of $G \setminus F_2$ and apply Lemma 6.16. We then see that G has a clique cutset, so conclusion (c) holds.

7 Proof of Theorem 7.1

Theorem 7.1 Let $\ell \geq 3$ be an integer. If G is a graph in $C_{2\ell+1}$ then one of the following holds:

- (a) G is a ring of length $2\ell + 1$;
- (b) G is a proper blowup of a twinless odd ℓ -template;
- (c) G has a universal vertex or
- (d) G has a clique cutset.

Proof. By Lemma 3.10, G contains no prism no theta and no proper wheel. Also, clearly G contains no C_4 and no C_5 . Hence, by Theorem 3.13, we may assume that G contains a pyramid for otherwise one of the conclusions (a), (c) or (d) holds. The result then follows from Lemma 6.1.

8 Even templates

8.1 Even ℓ -template partitions

For an integer $\ell \ge 4$, an even ℓ -template partition of a graph G is a partition of the vertex-set of G into five sets A, B, A', B' and I satisfying the following conditions.

- (a) $A = A_K \cup A_S$ where $A_K = \{v_1, \dots, v_k\}, A_S = \{v_{k+1}, \dots, v_{k+s}\}$ and $k+s \ge 3$.
- (b) $A' = A'_K \cup A'_S$ where $A'_K = \{v'_1, \dots, v'_k\}$ and $A'_S = \{v'_{k+1}, \dots, v'_{k+s}\}.$
- (c) For each $i \in \{1, \ldots, k\}$, v_i and v'_i are linked by a path of G of length $\ell 1$ and for each $i \in \{1, \ldots, s\}$, v_{k+i} and v'_{k+i} are linked by a path of G of length $\ell 2$. These k + s paths are vertex disjoint and they are called the *principal paths* of the partition.
- (d) I is the set of all internal vertices of the principal paths, every vertex in I has degree 2 in G.
- (e) Both A_K and A'_K are cliques of G and both A_S and A'_S are stable sets of G. For $i \in \{1, \ldots, k\}$ and $j \in \{1, \ldots, s\}$, exactly one of $v_i v_{k+j}$ and $v'_i v'_{k+j}$ is an edge. Furthermore G[A] and hence G[A'] are threshold graphs.
- (f) There exists a laminar hypergraph \mathcal{H} on vertex set $\{v_1, \ldots, v_{k+s}\}$ such that:
 - every hyperedge X of \mathcal{H} is an anticonnected module of G[A] of cardinality at least 2 and
 - if G[A] is not connected then at least one hyperedge of \mathcal{H} contains all vertices of A.
- (g) There exists a laminar hypergraph \mathcal{H}' on vertex set $\{v'_1, \ldots, v'_{k+s}\}$ such that:

- every hyperedge X' of \mathcal{H}' is a module of G[A'] of cardinality at least 2 and
- if G[A'] is not connected then least one hyperedge of \mathcal{H}' contains all vertices of A'.
- (h) $B = \{v_X : X \text{ hyperedge of } \mathcal{H}\}, B' = \{v'_X : X \text{ hyperedge of } \mathcal{H}'\}.$ The set of edges of G incident to vertices in $B \cup B'$ is defined as follows:
- (i) for every $v_X, v_Y \in B, v_X v_Y \in E(G)$ if and only if $X \cap Y \neq \emptyset$,
- (j) for every $v'_X, v'_Y \in B', v'_X v'_Y \in E(G)$ if and only if $X \cap Y \neq \emptyset$,
- (k) for every $v_i \in A$, $v_X \in B$, $v_i v_X \in E(G)$ if and only if $v_i \in N_{G[A]}[X]$,
- (l) for every $v'_i \in A', v'_X \in B', v'_iv'_X \in E(G)$ if and only if $v'_i \in N_{G[A']}[X]$.

The following notation is convenient.

Notation: For every vertex $x \in B$ such that $x = v_X$ where X is a hyperedge of \mathcal{H} , we set $H_x = X$. Similarly, for every vertex $x \in B'$ such that $x = v'_X$ where X is a hyperedge of \mathcal{H}' , we set $H'_x = X$.

We now list some properties of even ℓ -template partitions that follow directly from the definition.

- (i) G[A] and G[A'] are threshold graphs such that, $-A = A_K \cup A_S$ and $A' = A'_K \cup A'_S$, $-G[A_K]$ and $G[A'_K]$ are complete graphs having the same number of vertices, $-G[A_S]$ and $G[A'_S]$ are complement of complete graphs having the same number of vertices, the subgraph of G[A] induced by the edges between A_K and A_S is isomorphic to the complement of the subgraph of G[A'] induced by the edges between A'_K and A'_S .
- (ii) For all $x \in B$, H_x is a module of G[A] and $G[H_x]$ is anticonnected. Also for all $x \in B'$, H'_x is a module of G[A'] and $G[H'_x]$ is anticonnected.
- (iii) G[B] is isomorphic to the line graph of the hypergraph \mathcal{H} on vertex set A and hyperedge set $\{H_x : x \in B\}$. Also G[B'] is isomorphic to the line graph of the hypergraph \mathcal{H}' on vertex set A' and hyperedge set $\{H'_x : x \in B'\}$. Hence G[B] and G[B'] are quasi-threshold graphs by Theorem 3.4.
- (iv) There is an edge between $v_i \in A$ and $x \in B$ if and only if $v_i \in N_A[H_x]$, and there is an edge between $v'_i \in A'$ and $x \in B'$ if and only if $v'_i \in N_{A'}[H'_x]$.

By the fact that G[A] and G[A'] are threshold graphs in even and odd template partitions, by Properties (ii) and (iv) of even template partitions (which are the same as Properties (ii) and (iv) of odd templates) and conditions (f) and (g) of even template partitions, we have the following lemma whose proof is similar to the one of Lemma 4.4.

Lemma 8.1 There exist vertices w and w' that are universal vertices in respectively $G[A \cup B]$ and $G[A' \cup B']$.

Lemma 8.2 For an integer $\ell \geq 4$, every theta Θ such that $\Theta \in C_{2\ell}$ has an even ℓ -template partition, every prism Σ such that $\Sigma \in C_{2\ell}$ has an even ℓ -template partition.

Proof. Since $\Theta \in C_{2\ell}$, its three paths have length ℓ . Let x and y be the common two extremities of these paths, let A and A' be respectively the set of neighbors of x and y and let I be the set of vertices of Θ that are not in $A \cup A' \cup \{x, y\}$. It is easy to verify that $(A, \{x\}, A', \{y\}, I, x, y)$ is an even ℓ -template partition. Similarly all three paths of Σ have length $\ell - 1$. Let A be the set of the vertices v_1, v_2, v_3 of one of the triangles and A' be the set of vertices v'_1, v'_2, v'_3 of the other triangle. It is easy to verify that $(A, \emptyset, A', \emptyset, V(\Sigma) \setminus (A \cup A'), v_1, v'_1)$ is an even ℓ -template partition. \Box

Due to the similarities in odd and even template partitions as pointed out above we also get the following results similar to Lemmas 4.6, 4.7 and 4.8 with the same proofs.

Lemma 8.3 If $x \in B$ (resp. $x \in B'$), then H_x (resp. H'_x) is the unique anticomponent of $G[N_A(x)]$ (resp. $G[N_{A'}(x)]$) that contains at least two vertices.

Lemma 8.4 If $x, y \in B$ (resp. $x, y \in B'$) are such that $xy \notin E(G)$, then $H_x \cup \{x\}$ (resp. $H'_x \cup \{x\}$) is anticomplete to $H_y \cup \{y\}$ (resp. $H'_y \cup \{y\}$).

Lemma 8.5 Every vertex of G has degree at least 2 and every vertex of $B \cup B'$ has degree at least 3.

We may also extend \mathcal{H} into a hypergraph \mathcal{H}_A with vertex-set A by adding to its edge-set the hyperedge $H_v = N_A[v] \cap \{u \in A : u \leq_{G[A]} v\}$ for every vertex $v \in A$. Similarly we extend \mathcal{H}' into a hypergraph $\mathcal{H}'_{A'}$. The following lemma has the same proof as Lemma 4.10. **Lemma 8.6** \mathcal{H}_A is a laminar hypergraph and $G[A \cup B]$ is isomorphic to its line graph (in particular, $G[A \cup B]$ is a quasi-threshold graph and therefore a chordal graph). A similar statements holds for \mathcal{H}'_A , and $G[A' \cup B']$.

8.2 Even ℓ -templates

We will now need more notion and notation.

Given an even ℓ -template partition of G, we define a hypergraph \mathcal{H}_G whose vertex set is $\{k+1, k+2, \ldots, k+s\}$ and whose hyperedges are sets of indices of the vertices of $A_S \cup A_{S'}$ in hyperedges of $\mathcal{H}_A \cup \mathcal{H}_{A'}$. More formally, $E(\mathcal{H}_G) = E_A \cup E_{A'}$ where

- $E_A = \{\{i : v_i \in H \cap A_S\} : H \text{ hyperedge of } \mathcal{H}_A \text{ s.t. } H \cap A_S \neq \emptyset\}$

 $-E_{A'} = \{\{i : v_i \in H \cap A'_S\} : H \text{ hyperedge of } \mathcal{H}_{A'} \text{ s.t. } H \cap A'_S \neq \emptyset\}.$

Notice that \mathcal{H}_G may contain distinct hyperedges containing the same set of vertices.

A circular sequence $C = (j_1, e_1, j_2, ..., j_t, e_t, j_1)$, where the j_i 's are distinct vertices of \mathcal{H}_G and the e_i 's are distinct hyperedges of \mathcal{H}_G , is said to be a hyper cycle of length t of \mathcal{H}_G if

- each j_i belongs to e_{i-1} and e_i (where $e_{t+1} = e_1$) and to no other hyperedge of C,
- any two distinct hyperedges of C that belong both to E_A or both to $E_{A'}$ are disjoint.

We notice that by definition each e_i contains j_i and j_{i+1} and no other vertex of \mathcal{C} , hence any two consecutive hyperedges e_i and e_{i+1} of \mathcal{C} have a non empty intersection and none is included in the other. So, since \mathcal{H}_A and $\mathcal{H}_{A'}$ are both laminar hypergraphs, the hyperedges of \mathcal{C} belong alternately to E_A and $E_{A'}$. In particular the length of \mathcal{C} is even.

For an integer $\ell \geq 3$ and a graph G, a strong even ℓ -template partition of G is an even ℓ -template partition (A, A', B, B', I) of G, such that \mathcal{H}_G contains no hyper cycle of length greater than 2.

A graph G which has a strong even ℓ -template partition is called an *even* ℓ -template. We will denote by (A, B, A', B', I, w, w') an even ℓ -template G with strong even ℓ -template partition (A, B, A', B', I) such that w is a universal vertex of $G[A \cup B]$ and w' is a universal vertex of $G[A' \cup B']$ (by Lemma 8.1, such w and w' do exist).

Lemma 8.7 Let G be an even ℓ -template and (A, A', B, B', I) be a strong even ℓ -template partition of G. Every hole H of G contains two principal paths of G and either

- these two principal paths have length $\ell 1$ and they induce H,
- one principal path has length $\ell 1$, the other has length $\ell 2$ and H contains exactly one more vertex which belongs to $A_K \cup B \cup A'_K \cup B'$,
- both principal paths have length $\ell 2$, H contains exactly two more vertices, one in $A_K \cup B$ and the other in $A'_K \cup B'$.

In all cases H has length 2ℓ , and so $G \in \mathcal{C}_{2\ell}$.

Proof. Let H be a hole of G. By Lemma 8.6, $G[A \cup B]$ contains no P_4 and no C_4 , so H cannot contain only vertices of $A \cup B$, and similarly, it cannot contain only vertices of $A' \cup B'$. So H must contain vertices of some principal path, and hence it contains an even number of principal paths.

In case H contains two principal paths of length $\ell - 1$ then their endpoints are linked by two disjoint edges (by the definition of an even template partition). So these paths form a hole of length 2ℓ and H contains no other principal path.

Assume now that H contains exactly one path $P_u = u \dots u'$ of length $\ell - 1$ and at least one path $P_v = v \dots v'$ of length $\ell - 2$, for some $u \in A_K$ and $v \in A_S$. By the definition of an even template partition, there exists exactly one edge between P_u and any principal path of length $\ell - 2$, hence H cannot contain three such paths and should hence contain exactly one, namely P_v . Up to symmetry, we may assume that $uv \notin E(G)$ and then the hole H is made of the path $uP_uu'v'P_vv$ of length $2\ell - 2$ and a path $Q = u \dots v$ of $G[A \cup B]$. By Lemma 8.6, Q has length at most 2 and since $uv \notin E(G)$, we get that Q = ubv for some $b \in A \cup B$. So H has length 2ℓ and since A_S is a stable set in G[A] we have $b \in A_K \cup B$ as claimed.

It remains to consider the case where all principal paths contained in H have length $\ell-2$. Assume that H contains $t \geq 3$ principal paths P_1, P_2, \ldots, P_t associated to some $v_{j_1}, v_{j_2}, \ldots, v_{j_t} \in A_S$ (note that since t is even, then $t \geq 4$). By the definition of an even template partition, there exists no edge connecting these paths. Since by Lemma 8.6 there is no P_4 in $G[A \cup B]$ and in $G[A' \cup B']$, without loss of generality we may assume that $H = v_{j_1}P_1v'_{j_1}b_1v'_{j_2}P_2v_{j_2}b_2v_{j_3}, \ldots, v'_{j_t}P_tv_{j_t}b_tv_{j_1}$ where b_i 's with odd index belong to $A'_K \cup B'$ and those with even index belong to $A_K \cup B$. Hence, to each b_i we may associate the hyperedge e_i of \mathcal{H}_G corresponding to $H_{b_i} \in \mathcal{H}_A \cup \mathcal{H}_{A'}$. We claim that $\mathcal{C} = (j_1, e_1, j_2, \ldots, j_t, e_t, j_1)$ is a hyper cycle of \mathcal{H}_G . Assume there

exists in \mathcal{C} an hyperedge $e_l \neq e_{i-1}, e_i$ such that $j_i \in e_l$. So, up to symmetry, $v_{j_i} \in H_{b_l}$ and then by the definition of an even template, H would contain a chord $v_{j_i}b_l$, a contradiction. So the first condition for being a hyper cycle is satisfied by \mathcal{C} . Assume now, up to symmetry, that there exist $e_i, e_l \in H_A$ that are not disjoint. Then by the definition of an even template partition, H would contain a chord $b_i b_l$, a contradiction again. Hence the second condition should be satisfied and \mathcal{C} is a hyper cycle of \mathcal{H}_G , a contradiction to the fact that (A, A', B, B', I) is a strong even ℓ -template partition of a graph G. So, we may now conclude that H contains exactly two principal paths of length $\ell - 2$ and two more vertices $b_1 \in A_K \cup B$ and $b_2 \in A'_K \cup B'$.

8.3 Even ℓ -pretemplates

For every integer $\ell \geq 4$, an even ℓ -pretemplate partition of a graph G is a partition of the vertex-set of G into five sets $A = A_K \cup A_S$, $B, A' = A'_K \cup A'_S$, B' and I that satisfy the following conditions.

- 1. $N(B) \subseteq A$ and $N(A \cup B) \subseteq I$.
- 2. $N(B') \subseteq A'$ and $N(A' \cup B') \subseteq I$.
- 3. $|A_K| = |A'_K| = k, A_K = \{v_1, \dots, v_k\}$ and $A'_K = \{v'_1, \dots, v'_k\}.$
- 4. $|A_S| = |A'_S| = s$, $A_S = \{v_{k+1}, \dots, v_{k+s}\}$ and $A'_S = \{v'_{k+1}, \dots, v'_{k+s}\}$ are stable sets of G where $k + s \ge 3$.
- 5. For every $i \in \{1, \ldots, k+s\}$, there exists a unique path P_i from v_i to v'_i whose interior is in I.
- 6. Every vertex in I has degree 2 and lies on a path from v_i to v'_i for some $i \in \{1, \ldots, k\}$.
- 7. All paths P_1, \ldots, P_k have length $\ell 1$, all paths P_{k+1}, \ldots, P_{k+s} have length $\ell 2$.
- 8. $G[A \cup B]$ and $G[A' \cup B']$ are both connected graphs.
- 9. Every vertex of B is in the interior of a path of $G[A \cup B]$ with both ends in A.
- 10. Every vertex of B' is in the interior of a path of $G[A' \cup B']$ with both ends in A'.

We then say that (A, A', B, B', I) is an even ℓ -pretemplate partition of G.

Lemma 8.8 Let $\ell \geq 4$ be an integer and G be a graph of $C_{2\ell}$. Any even ℓ -pretemplate partition (A, B, A', B', I) of G is a strong even ℓ -template partition of G.

Proof.

(1) For all distinct $i, j \in \{1, ..., k\}$, $v_i v_j \in E(G)$ and $v'_i v'_j \in E(G)$. In particular, A_K and A'_K are cliques of G.

Assume that at least one of $v_i v_j, v'_i v'_j \notin E(G)$. By condition 8. in the definition of an even pretemplate partition there exist a path between v_i and v_j in $G[A \cup B]$ and a path between v'_i and v'_j in $G[A' \cup B']$. Together with P_i and P_j , these paths form a hole of length at least $2\ell + 1$, a contradiction. This proves (1).

(2) For all distinct $i \in \{1, ..., k\}$ and $j \in \{k + 1, ..., k + s\}$, exactly one of $v_i v_j$ and $v'_i v'_j$ is an edge of G.

By condition 8. in the definition of an even pretemplate partition there exist a path between v_i and v_j in $G[A \cup B]$ and a path between v'_i and v'_j in $G[A' \cup B']$. Together with the principal paths P_i (of length $\ell - 1$) and P_j (of length $\ell - 2$), these paths form a hole which will be of length 2ℓ if and only if one of the paths is of length 1 and the other path is of length 2. This proves (2).

(3) Every path of $G[A \cup B]$ with both ends in A is of length at most 2. The same holds for a path of $G[A' \cup B']$ with both ends in A'.

Assume on the contrary that there exists a path P of length at least 3 in $G[A \cup B]$ with both ends v_i, v_j in A. Then by (1) and condition 7. of an even pretemplate partition, at least one of P_i, P_j is of length $\ell - 2$, say P_j . If P_i is of length $\ell - 1$ then by (2) $v'_i v'_j \in E(G)$ and P, P_i, P_j would induce a hole of length at least $2\ell + 1$, a contradiction. Hence P_i and P_j are both of length $\ell - 2$ and by conditions 4. and 8. of an even pretemplate partition, P, P_i, P_j and any path between v'_i and v'_j form a hole of length at least $2\ell + 1$, a contradiction again. The proof for $G[A' \cup B']$ is similar. This proves (3).

(4) G[A] is a threshold graph. The same holds for G[A'].

G[A] is obviously C_4 -free. By (1) and condition 4. of an even pretemplate partition, A is partitioned into a clique and a stable set of G, so by Theorem 3.3 G[A] is $2K_2$ -free. By (3), G[A] is P_4 -free. So G[A] is $(P_4, C_4, 2K_2)$ -free and is therefore a threshold graph. The proof for G[A'] is similar. This proves (4).

We now study the structure of G[B] (respectively G[B']) and its relation with G[A] (respectively G[A']).

(5) For every vertex $x \in B$, $G[N_A(x)]$ has a unique anticonnected component of size at least 2. The same holds for $G[N_{A'}(x)]$ when x is any vertex in B'.

By condition 9. of an even pretemplate partition, x is in the interior of a path $P = v_i \dots v_j$ of $G[A \cup B]$ with both ends in A. By (3), P has length 2, so x is adjacent to v_i and v_j . Hence $G[N_A(x)]$ has an anticonnected component of size at least 2. It is unique, for otherwise G[A] would contain a C_4 . This proves (5).

For all $x \in B$ (respectively B'), we define H_x to be the anticonnected component of $G[N_A(x)]$ (respectively $G[N'_A(x)]$) of size at least 2 whose existence follows from (5).

(6) For every x in B (respectively B'), H_x is a module of G[A] (respectively G[A']).

Let $x \in B$. If the claim does not hold, since H_x is by definition anticonnected, there exist $v_h \in A \setminus H_x$ and non-adjacent $v_i, v_j \in H_x$ such that $v_h v_i \in E(G)$ and $v_h v_j \notin E(G)$. Note that $xv_h \notin E(G)$ because otherwise, v_h would be in H_x . Then $v_h v_i x v_j$ is a path of length 3, a contradiction to (3). The similar proof holds for x in B'. This proves (6).

(7) If xy is an edge of G[B] or G[B'], then $H_x \subseteq H_y$ or $H_y \subseteq H_x$.

Let xy be an edge of G[B]. Up to symmetry, we may assume that $N_A(x) \subseteq N_A(y)$, for otherwise vertices $v_i \in N_A(x) \setminus N_A(y)$ and $v_j \in N_A(y) \setminus N_A(x)$ either form a C_4 with x and y or a contradiction to (3).

By (5), $G[N_A(y)]$ has only one anticonnected component of size at least 2, namely H_y . Since H_x is anticonnected, has size at least 2 and is included in $N_A(y)$, it must be included in H_y . The similar proof holds for an edge xyof G[B']. This proves (7).

(8) If x and y are non-adjacent vertices of B or B' then H_x and H_y are disjoint.

On the contrary, suppose that x and y are nonadjacent vertices of B but there exists a vertex $v \in H_x \cap H_y$. Since H_x is anticonnected and of size at least 2, there exists $v_i \in H_x$ non-adjacent to v. Note that $v_i y \notin E(G)$, for otherwise x, y, v_i and v form a C_4 . Similarly, there exists a vertex $v_i \in H_y$ that is non-adjacent to v and to x. If $v_i v_j \in E(G)$, then $\{x, y, v, v_i, v_j\}$ induces a C_5 , a contradiction. Otherwise, $\{x, y, v, v_i, v_j\}$ induces a P_5 , a contradiction to (3). The proof for $x, y \in B'$ is similar. This proves (8).

We are now ready to define the hypergraphs \mathcal{H} and \mathcal{H}' . For every $x \in B$ (respectively B'), we defined a set $H_x \subseteq A$ (respectively $H_x \subseteq A'$). From (7) and (8), the sets H_x for $x \in B$ form a laminar hypergraph \mathcal{H} (with vertex set A). Symetrically, the sets H_x for $x \in B'$ form a laminar hypergraph \mathcal{H}' (with vertex set A').

(9) If G[A] is not connected then at least one hyperedge of \mathcal{H} contains all vertices of A.

Assume G[A] is not connected. By (4), G[A] is a threshold graph, and then by Theorem 3.7 it contains an isolated vertex v_i . By the definition of an even pretemplate partition, $G[A \cup B]$ is connected and $|A| \ge 3$, so there exists a path P in $G[A \cup B]$ from v_i to a vertex $u \in A \setminus \{v_i\}$. By (3) and since v_i has no neighbor in A, we have that $P = uyv_i$ where $y \in B$. So, H_y contains v_i . We may therefore consider the hyperedge W of \mathcal{H} that contains v_i and that is inclusion-wise maximal w.r.t. this property. If there exists $v_j \in A \setminus W$, since $v_j v_i \notin E(G)$, we deduce as above that \mathcal{H} has a hyperedge Z that contains i and j. Because of v_j , $Z \subseteq W$ is impossible; because of v_i , $W \cap Z = \emptyset$ is impossible; and because of the maximality of $W, W \subsetneq Z$ is impossible. Hence, W and Z contradict \mathcal{H} being laminar. So W = A.

This proves (9).

At this point we can easily verify that (A, B, A', B', I) is an even ℓ -template partition of G:

- conditions (a) (b) (c) and (d) of an even template partition are satisfied because of conditions 3. 4. 5. 6. and 7. in the definition of an even pretemplate partition,

- (e) of an even template partition is a consequence of condition 4. of an even pretemplate partition and (1), (2), (4),

- (f), (g) and (h) of even template partition is a consequence of (5), (6), (7), (8),

- (i) and (j) of an even template partition comes from (7) and (8),

- (k) and (l) of an even template partition follows easily from our previous results : By definition of H_x , for every x in B, $N_A(x) \subseteq N_A[H_x]$. Suppose that there exists $u \in N_A[H_x] \setminus N_A(x)$. Since by (6) H_x is a module, it follows from Lemma 4.2 that u is complete to H_x , so x and u together with two non-adjacent vertices from H_x induce a C_4 , a contradiction. Hence, $N_A(x) = N_A[H_x]$ and condition (k) of an even template partition is satisfied. The proof for (1) is similar.

It remains to prove the following.

(10) \mathcal{H}_G contains no hyper cycle of length greater than 2.

Suppose on the contrary that $\mathcal{C} = (j_1, e_1, j_2, ..., j_t, e_t, j_1)$ is a hyper cycle of \mathcal{H}_G of length t > 2. Without loss of generality, each e_i with odd index belongs to E_A , and since it contains j_i and j_{i+1} such that $v_{j_i}, v_{j_{i+1}} \in A_S$ we have that e_i is the set of indices of the vertices of H_{b_i} for some $b_i \in A_K \cup B$. Similarly each e_i with even index belongs to $E_{A'}$, contains j_i and j_{i+1} such that v'_{j_i} and $v'_{j_{i+1}} \in A'_S$ and corresponds to $H_{b'_i}$ for some $b'_i \in A'_K \cup B'$. Hence $C = v_{j_1} b_1 v_{j_2} P_{v_{j_2}} v'_{j_2} b'_2 v'_{j_3} P_{v_{j_3}} v_{j_3} \dots v'_{j_t} b'_t v'_{j_1} P_{v_{j_1}} v_{j_1}$ is a cycle in G of length $t\ell > 2\ell$.

By definition of an even template partition, there are no edges between two distinct P_i 's contained in C, and no edges between vertices in $A_K \cup B$ and vertices in $A'_K \cup B'$. So the only chords C could contain are:

- $b_i b_j$ for some distinct $i, j \in \{1, 2, \dots, t\}$ of same parity, or
- $v_{j_l}b_i$ for some odd $i \in \{1, 2, \ldots, t\}$ and $l \neq i, i+1$ (where t+1=1), or $v'_{j_l}b_i$ for some even $i \in \{1, 2, \ldots, t\}$ and $l \neq i, i+1$ (where t+1=1).

The first case is impossible since, by definition of a hyper cycle, the hyperedges H_{b_i} are all disjoint. The two other cases cannot occur because else j_l would be contained in the hyperedge e_i where $i \neq j_{l-1}, j_l$, a contradiction to the definition of a hyper cycle.

So C is a hole which has length more than 2ℓ , a contradiction to $G \in \mathcal{C}_{2\ell}$. This proves (10).

This ends the proof of Lemma 8.8.

Notice that as a corollary of Lemma 8.8 we get that if a graph G in $\mathcal{C}_{2\ell}$ has an even ℓ -pretemplate partition then it is an even ℓ -template and any of its even ℓ -pretemplate partition is a strong even ℓ -template partition.

We will need the following analog of Lemma 4.15.

Lemma 8.9 Let G = (A, B, A', B', I, w, w') be an even ℓ -template. Two vertices x and y of G are twins if and only if $x, y \in B$ and $H_x = H_y$, or $x, y \in B'$ and $H'_x = H'_y$.

Proof. If $x, y \in B$ and $H_x = H_y$, or $x, y \in B'$ and $H'_x = H'_y$, then x and y are obviously twins.

We claim that for all $x \in A \cup I \cup A'$, there exist two vertices $a, b \in N_G(x)$ such that $N[a] \cap N[b] = \{x\}$. If $x \in I$, choose a and b to be the only two neighbors of x. If $x \in A$, then let a be the neighbor of x in I and let b = w if $x \neq w$, else let b be any vertex of $A \setminus \{x\}$ (we remind that w is by definition a vertex universal in $G[A \cup B]$). In all cases, by definition of an even template partition, $N_G[a] \cap N_G[b] = \{x\}$. The proof is similar when $x \in A'$. So, x has no twin in G.

An even ℓ -partition (A, B, A', B', I) of an even ℓ -template G is proper if all universal vertices of $G[A \cup B]$ (respectively $G[A' \cup B']$) are in B (respectively B').

Lemma 8.10 Every twinless even ℓ -template G = (A, B, A', B', I, w, w')with a proper even ℓ -partition satisfies the following:

- $w \in B$, w is the unique universal vertex of $G[A \cup B]$ and G[A] contains at least one isolated vertex $u \in A_S$,
- $w' \in B'$, w' is the unique universal vertex of $G[A' \cup B']$ and G[A']contains at least one isolated vertex $v' \in A'_S$,
- $|A_S| \ge 2.$

Proof. Since the partition is proper, all universal vertices of $G[A \cup B]$ belong to B and since G is twinless, B contains a unique vertex which is universal in $G[A \cup B]$. Hence w, which is by definition of G a universal vertex of $G[A \cup B]$, belongs to B and G[A] contains no universal vertex since else, by the definition of an even template, this vertex would be universal in $G[A \cup B]$. Then, by Theorem 3.4, G[A] contains at least one isolated vertex say u. The proof for w' is similar and there exists a vertex v' isolated in G[A']. It remains to prove the third assertion. Assume that $|A_S| \leq 1$. Then, since A contains at least three vertices, it should be that $A_S = \{u\}$. Similarly we obtain that $A'_S = \{v'\}$, so u' = v'. Then by the definition of an even template v' is not isolated in G[A'], a contradiction. \Box

Lemma 8.11 For every integer $\ell \geq 4$, every even ℓ -template G admits an even ℓ -partition which is proper.

Proof. Let G = (A, B, A', B', I) be a strong even ℓ -partition of G. By Lemma 8.1 we know that each of $G[A \cup B]$ and $G[A' \cup B']$ contains at least one universal vertex. If the lemma does not hold, we may assume up to symmetry, that there exists at least one vertex w universal in $G[A \cup B]$ which is in A. Then from the definition of an even template it is clear that we may choose w in A_K . We denote by w_+ the neighbor of w on the principal path P_w of G in the partition (A, B, A', B', I). Since $\ell \geq 4$, we have $w_+ \in I$. Consider now the partition $(\mathbb{A}_K, \mathbb{A}_S, \mathbb{B}, \mathbb{A}'_K, \mathbb{A}'_S, \mathbb{B}', \mathbb{I})$ of V(G) where $\mathbb{A}_K = A_K \setminus \{w\}$, $\mathbb{A}_S = A_S \cup \{w_+\}, \mathbb{B} = B \cup \{w\}, \mathbb{A}'_K = A'_K \setminus \{w'\}, \mathbb{A}'_S = A'_S \cup \{w'\}, \mathbb{B}' = B',$ $\mathbb{I} = I \setminus \{w_+\}$. We set $\mathbb{A} = \mathbb{A}_K \cup \mathbb{A}_S$ and $\mathbb{A}' = \mathbb{A}'_K \cup \mathbb{A}'_S$.

Since (A, B, A', B', I) is an even ℓ -partition of G, it is clear that A_K and \mathbb{A}'_K are cliques of the same cardinality. Since the only edge in G between w_+ and A is ww_+ we have that $\mathbb{A}_S = A_S \cup \{w_+\}$ is a stable set and since w_+ has no neighbor in \mathbb{A} , any universal vertex of $G[\mathbb{A} \cup \mathbb{B}]$ is in \mathbb{B} . As w is complete to A_S we get from the definition of an even template that $\mathbb{A}'_S = A'_S \cup \{w'\}$ is a stable set of the same cardinality as $A_S \cup \{w_+\}$. Remark that $|\mathbb{A}| = |A| \ge 3$. Furthermore there exists a unique path $P_{w_{+}} = w_{+}P_{w}w'$ between w_{+} and w'whose interior is in I. This path has length $\ell - 2$. It is also important to notice that w is in the interior of a path vww_+ for any $v \in A \setminus \{w\}$. Every other vertex of \mathbb{B} is in the interior of a path of $G[\mathbb{A} \cup \mathbb{B}]$ with both ends in \mathbb{A} , path which is the same as in the initial partition since obviously this path did not contain w. With all these observations it is easy to conclude that $\mathbb{P} = (\mathbb{A}, \mathbb{B}, \mathbb{A}', \mathbb{B}', \mathbb{I})$ fulfills all conditions to be a pretemplate partition of G. By Lemma 8.7 we know that $G \in \mathcal{C}_{2\ell}$. So by Lemma 8.8 and the fact that w_+ is isolated in $G[\mathbb{A}], \mathbb{P}$ is an even ℓ -partition of G such that no vertex of A is universal in $G[\mathbb{A} \cup \mathbb{B}]$.

If \mathbb{A}' contains no universal vertex of $G[\mathbb{A}' \cup \mathbb{B}']$ then \mathbb{P} is a proper even ℓ -partition of G, else a proof similar to the one above allows to obtain from \mathbb{P} a proper even ℓ -partition of G.

Lemma 8.12 For every integer $\ell \geq 4$, every even ℓ -template G contains a prism or a theta.

Proof. By Lemma 8.11, G admits a proper even ℓ -partition (A, B, A', B') of G and by definition of an even template, G contains three vertices v_1, v_2 and v_3 in A and the corresponding vertices v'_1, v'_2 and v'_3 in A'. Considering all kinds of repartition of v_1, v_2 and v_3 in A_K and A_S , all kinds of attachment between them respecting the rules of the partition and the fact that B (respectively B') contains a vertex universal in $G[A_K \cup A_S \cup B]$ (respectively $G[A'_K \cup A'_S \cup B']$) it is easy to verify that in each case we obtain either a prism or a theta.

8.4 Blowups and holes

We may define flat, solid and optional edges in even ℓ -template partitions exactly as in the case of odd ℓ -template partitions (see subsection 5.1). In the following we will also use the notion of blowup and preblowup of an even template ℓ -partition with the same definition as for an odd ℓ -template partition (see subsections 5.2 and 5.3).

The following lemma can be proved similarly as Lemma 5.4.

Lemma 8.13 A hole C in a blowup of a twinless even ℓ -template contains at most one vertex in each blown up clique.

We remark that Lemmas 5.1, 5.2 and 5.3 are valid for any odd or even template as they rely only on the definition of solid and optional edges. Hence, the following lemma has the same proof as Lemma 5.5 of the odd case, except that we use Lemma 8.7 instead of Lemma 4.11, and Lemma 8.13 instead of Lemma 5.4.

Lemma 8.14 In a blowup G^* of a twinless even ℓ -template G, every hole has length 2ℓ .

Recall that to blowup (resp. preblowup) a template, one needs to first fix an ℓ -partition. If this partition is proper, the blowup (resp. preblowup) is *proper*. Recall also that when G^* is a preblowup of a template G, the *domination score* of G w.r.t. G^* is defined as (where N refers to the neighborhood in G^*):

$$s(G, G^*) = \sum_{x \in A \cup A' \cup I} |\{x^* \in K_x : N[x^*] \subseteq N[x]\}|$$

Lemma 8.15 Let $\ell \geq 3$ and let G^* be a proper preblowup of an even ℓ -template with $k \geq 3$ principal paths. If $G^* \in C_{2\ell}$, then G^* is a proper blowup of a twinless even ℓ -template G with k principal paths (in particular, G is an induced subgraph of G^*).

Proof. Among all the induced subgraphs of G^* that are even ℓ -templates and for which G^* is a proper preblowup, we suppose that G is one that maximizes $s(G, G^*)$. We denote by (A, B, A', B', I, w, w') the proper ℓ -partition of G that is used for its preblowup and by $(A^*, B^*, A'^*, B'^*, I^*)$ the corresponding partition of the vertices of G^* .

(1) There exist vertices $w^* \in B^*$ and $w'^* \in B'^*$ that are complete to respectively A^* and A'^* .

Since the partition is proper, $w \in B$ and then by condition (b1) of a preblowup, there exists $w^* \in B^*$ that is complete to A^* .

The proof of the statement about $w^{\prime*}$ is similar. This proves (1).

(2) For every principal path $P_u = u \dots u'$ of G and $u^* \in K_u$, there exists in G^* a path P_{u^*} from u^* to some $u'^* \in K_{u'}$ whose interior is in $\bigcup_{x \in I \cap V(P_u)} K_x$ and whose length is equal to the length of P_u . Moreover, the interior of P_{u^*} is anticomplete to $V(G^*) \setminus \bigcup_{v \in V(P_u)} K_v$.

The existence of a path of same length as P_u from u^* to some $u'^* \in K_{u'}$ whose interior is in $\bigcup_{x \in I \cap V(P)} K_x$ follows from conditions (a2), (i) and (i1), of preblowup. The statement about its interior follows from conditions (a), (b) and (i) of preblowup. This proves (2).

(3) For all $u, v \in A$ such that $uv \notin E(G)$, K_u is anticomplete to K_v . A similar statement holds for A'.

Suppose that there exist $u^* \in K_u$ and $v^* \in K_v$ such that $u^*v^* \in E(G^*)$. By condition (a1) of preblowup, $u \neq u^*$ and $v \neq v^*$.

Since $uv \notin E(G)$ then at least one of u, v is in A_S , say $v \in A_S$.

Consider first the case where $u \in A_K$. The principal paths $P_u = u \dots u'$ and $P_v = v \dots v'$ have length respectively $\ell - 1$ and $\ell - 2$. Denote by u^+ the neighbor of u in P_u and by v^+ the neighbor of v in P_v . By property (e) of an even template, $u'v' \in E(G)$. Hence $uP_uu'v'P_vvv^*u^*u$ is a cycle C of length $2\ell + 1$. By conditions (a) and (a1) of preblowup, the only possible chords in C are u^*u^+ and v^*v^+ . Assume that $u^*u^+ \in E(G^*)$. Let P_{v^*} be a path of length $\ell - 2$ from v^* to v'^* as defined in (2). Since $v'^* \in K_{v'}$ and by (a1) of the preblowup applied to A', $v'^*u' \in E(G^*)$. So $v^*P_{v^*}v'^*u'P_uu^+u^*v^*$ is a hole of length $2\ell - 1$, a contradiction. Hence v^+v^* should be a chord of C. Let then P_{u^*} be a path of length $\ell - 1$ from u^* to u'^* as defined in (2). Since $u'^* \in K_{u'}$ and by (a1) of the preblowup applied to A', $u'^*v' \in E(G^*)$. So $u^*P_{u^*}u'^*v'P_vv^+v^*u^*$ is a hole of length $2\ell - 1$, a contradiction again.

It remains to consider the case where both u and v are in A_S . Let $P_u = u \dots u'$ and $P_v = v \dots v'$ be principal paths of length $\ell - 2$. Denote by u^+ the neighbor of u in P_u and by v^+ the neighbor of v in P_v . By property (e) of an even template, $u'v' \notin E(G)$, hence $uP_uu'w'^*v'P_vvv^*u^*u$ is a cycle C of length $2\ell + 1$. By conditions (a) and (a1) of preblowup, the only possible chords in C are u^+u^* and v^+v^* .

Assume without loss of generality that $u^+u^* \in E(G^*)$. Let P_{v^*} be a path of length $\ell - 2$ from v^* to v'^* as defined in (2). Since $v'^* \in K_{v'}$ and by (a1) applied to A', $v'^*u' \in E(G^*)$. So $v^*P_{v^*}v'^*w'^*u'P_uu^+u^*v^*$ is a hole of length $2\ell - 1$, a contradiction.

The result for A' holds symmetrically. This proves (3).

(4) For all $u, v \in A$ such that $uv \in E(G)$, K_u is complete to K_v . A similar statement holds for A'.

Suppose that there exist $u^* \in K_u$ and $v^* \in K_v$ such that $u^*v^* \notin E(G^*)$. Let $P_{u^*} = u^* \dots u'^*$ and $P_{v^*} = v^* \dots v'^*$ be defined as in (2). Observe that $u'^* \in K_{u'}$ and $v'^* \in K_{v'}$.

Since $uv \in E(G)$ then at least one of u, v is in A_K , say $u \in A_K$.

Consider first the case where $v \in A_K$ too. Then both P_{u^*} and P_{v^*} have length $\ell-1$. If $u'^*v'^* \in E(G^*)$ then $u^*P_{u^*}u'^*v'^*P_{v^*}v^*w^*u^*$ is a hole of length $2\ell+1$ and else $u^*P_{u^*}u'^*w'^*v'^*P_{v^*}v^*w^*u^*$ is a hole of length $2\ell+2$, so it should be that $v \in A_S$. Then P_{u^*} and P_{v^*} have length respectively $\ell-1$ and $\ell-2$. Moreover by property (e) of an even template, $u'v' \notin E(G^*)$, so by (3) $u'^*v'^* \notin E(G^*)$. Now $u^*P_{u^*}u'^*w'^*P_{v^*}v^*w^*u^*$ is a hole of length $2\ell+1$, a contradiction again.

This proves (4).

(5) For all $u \in I$ and $u_1, u_2 \in K_u$, either $N[u_1] \subseteq N[u_2]$ or $N[u_2] \subseteq N[u_1]$.

Otherwise, there exist $x_1^* \in N[u_1] \setminus N[u_2]$ and $x_2^* \in N[u_2] \setminus N[u_1]$. Note that $x_1^*x_2^* \notin E(G^*)$ for otherwise, $\{x_1^*, x_2^*, u_1, u_2\}$ induces a C_4 . It follows that x_1^* and x_2^* belong respectively to distinct cliques K_{x_1} and K_{x_2} , where x_1 and x_2 are the two neighbors of u along some principal path $P_v = v \dots v'$ of G. Because of x_1^*, x_2^* and condition (i1) of preblowup, there exists a path P^* from some $v^* \in K_v$ to some $v'^* \in K_{v'}$ whose interior is in $\bigcup_{x \in I \cap V(P_v)} K_x$ which contains u_1 and u_2 and has a length equal to the length of P_v plus one.

Assume first that $v \in A_K$. Then P^* has length ℓ . By Lemma 8.10, since (A, B, A', B', I, w, w') is a proper ℓ -partition of G, A_S is not empty. Let $q \neq v$ be a vertex in A_S and $P_q = q \dots q'$ be the principal path of length $\ell-2$ joining q and q' in G. Up to symmetry we may assume that $qv \notin E(G)$ and $q'v' \in E(G)$. Now, by condition (i) of preblowup and (1), P^* , P_q and w^* form a hole of length $2\ell + 1$, a contradiction.

So v should be in A_S and P^* has length $\ell - 1$. Since the ℓ -partition of G is proper, there exists a vertex $q \in A_S$ distinct from v, P_q has length $\ell - 2$ and there exists no edge between P^* and P_q . Then P^*, P_q, w^* and w'^* induce a hole of length $2\ell + 1$, a contradiction again. This proves (5).

(6) For all $u \in I$ and $u^* \in K_u$, $N[u^*] \subseteq N[u]$.

Otherwise, by (5), there exists a vertex $u^* \in K_u$ such that $N[u] \subsetneq N[u^*]$. Hence $(V(G) \setminus \{u\}) \cup \{u^*\}$ induces a subgraph G_0 of G^* and it is easy to verify that G^* is a preblowup of G_0 . This contradicts the maximality of $s(G, G^*)$. This proves (6).

By (5), for every $u \in I$, the clique K_u can be linearly ordered by the inclusion of the neighborhoods as u_1, \ldots, u_{k_u} with $u = u_{k_u}$ by (6) (so, for $1 \leq i \leq j \leq k_u$, $N[u_i] \subseteq N[u_j]$). From condition (i) of the preblowup it also follows that, in G^* , u is complete to the cliques associated to its two neighbors in G.

(7) For every $u \in A$ and $u_1, u_2 \in K_u$, either $N[u_1] \subseteq N[u_2]$ or $N[u_2] \subseteq N[u_1]$. A similar statement holds for A'.

Otherwise, there exist $x_1 \in N[u_1] \setminus N[u_2]$ and $x_2 \in N[u_2] \setminus N[u_1]$. Note that $x_1x_2 \notin E(G^*)$ for otherwise, $\{x_1, x_2, u_1, u_2\}$ induces a C_4 .

Observe first that by (3) and (4), $N_{A^*}[u_1] = N_{A^*}[u_2]$. Hence by condition (a) of preblowup, $x_1, x_2 \in B^* \cup K_{u^+}$ where u^+ is the neighbor of u in the principal path that contains u. Without loss of generality and since K_{u^+} is a clique, $x_1 \in B^*$.

By condition (b2) of preblowup, there exist non-adjacent $a, b \in A$ such that x_1 has neighbors $a^* \in K_a$ and $b^* \in K_b$, and by (3) $a^*b^* \notin E(G^*)$. Note that $a^*, b^* \neq u_2$ because $u_2x_1 \notin E(G^*)$. If u_2 is complete to $\{a^*, b^*\}$, then $\{u_2, a^*, x_1, b^*\}$ induces a C_4 , a contradiction. So, up to symmetry $u_2a^* \notin E(G)$. So, $a^* \notin K_u$ and by (4) and (3), $a^*u_1 \notin E(G^*)$. Observe that $x_2a^* \notin E(G^*)$ for otherwise $\{a^*, x_1, u_1, u_2, x_2\}$ induces a C_5 .

Suppose that $x_2 \in B^*$. As above, we can show that x_2 has a neighbor $c^* \in A^*$ that is anticomplete to $\{u_1, u_2, x_1\}$. Let $P_{a^*} = a^* \dots a'^*$ and $P_{c^*} = c^* \dots c'^*$ be defined as in (2). To avoid a hole $c^* x_2 u_2 u_1 x_1 a^* c^*$ of length 6, we have $a^* c^* \notin E(G^*)$. Since the lengths of P_{a^*} and P_{c^*} are at least $\ell - 2$, depending whether $a'^* c'^* \in E(G^*)$, $u_1 x_1 a^* P_{a^*} a'^* c'^* P_{c^*} c^* x_2 u_2 u_1$ is a hole of length at least $2\ell + 2$, a contradiction. So $x_2 \in K_{u^+}$. Using condition (i1) of preblowup, it is easy to verify that there exists a path P_{u_2} from u_2 to some $u'^* \in K_{u'}$ defined similarly than in (2), which contains x_2 . Now $u_2 P_{u_2} u'^* a'^* P_{a^*} a^* x_1 u_1 u_2$ or $u_2 P_{u_2} u'^* w'^* a'^* P_{a^*} a^* x_1 u_1 u_2$ is a hole of length $2\ell + 1$, a contradiction.

The result for A' holds symmetrically. This proves (7).

(8) For all $u \in A$ and $u^* \in K_u$, $N[u^*] \subseteq N[u]$. A similar statement holds for A'.

Otherwise, by (7) there exists a vertex $u^* \in K_u$ such that $N[u] \subsetneq N[u^*]$.
Hence, $(V(G) \setminus \{u\}) \cup \{u^*\}$ induces a subgraph G_0 of G^* which is a template (by Lemma 8.8) with a proper partition (by (3) and (4)). It is easy to verify that G^* is a preblowup of G_0 . This contradicts the maximality of $s(G, G^*)$. The result for A' holds symmetrically. This proves (8).

By (7), for every $u \in A \cup A'$, the clique K_u can be linearly ordered by the inclusion of the neighborhoods as u_1, \ldots, u_{k_u} , and by (8) $u_{k_u} = u$ (so, for $1 \leq i \leq j \leq k_u$, $N[u_i] \subseteq N[u_j]$).

(9) If xy is an edge of $G[B^*]$, then either $N_{A^*}(x) \subseteq N_{A^*}(y)$ or $N_{A^*}(y) \subseteq N_{A^*}(x)$.

Otherwise, there exist $u^* \in N_{A^*}(x) \setminus N_{A^*}(y)$ and $v^* \in N_{A^*}(y) \setminus N_{A^*}(x)$. Note that $u^*v^* \notin E$ for otherwise $\{u^*, x, y, v^*\}$ induces a C_4 . So, for some distinct $u, v \in A$, we have $u^* \in K_u$ and $v^* \in K_v$. Hence, by (4), $uv \notin E(G)$. Let $P_{u^*} = u^* \dots u'^*$ and $P_{v^*} = v^* \dots v'^*$ be defined as in (2). So, $xu^*P_{u^*}u'^*v'^*P_{v^*}v^*yx$ or $xu^*P_{u^*}u'^*v'^*P_{v^*}v^*yx$ forms a hole of length $2\ell+1$, a contradiction. This proves (9).

(10) For every $x \in B^*$, there exist non-adjacent $u, v \in A$ such that $xu, xv \in E(G^*)$.

This follows from condition (b2) of preblowup and from (8). This proves (10).

Two vertices x, y in B^* are equivalent if $N_A(x) = N_A(y)$.

(11) If x and y are equivalent vertices of B^* , then $xy \in E(G^*)$.

If $xy \notin E(G^*)$, then x, y and two of their neighbors provided by (10) induce a C_4 . This proves (11).

Vertices of B^* are partitioned into equivalence classes. By (11), each equivalence class is a clique X, and by (9), vertices of X can be linearly ordered according to the inclusion of neighborhoods in A^* . In each such a clique X we choose a vertex x maximal for the order and call B_1 the set of these maximal vertices. For every $x \in B_1$, we denote by K_x the clique of B^* of all vertices equivalent to x. Remind that $w^* \in B^*$: w^* is a maximal vertex of its clique. Hence, we can set $w^* \in B_1$.

So, for every $u \in B_1$, the clique K_u can be linearly ordered by the inclusion of the neighborhod in A^* as u_1, \ldots, u_{k_u} with $u = u_{k_u}$ (so, for $1 \le i \le j \le k_u, N_{A^*}(u_i) \subseteq N_{A^*}(u_j)$).

Statements similar to (9), (10), (11) hold for $B^{\prime*}$ and we define B_1^{\prime} as well.

We set $G_1 = G^*[A \cup B_1 \cup A' \cup B'_1 \cup I]$ and claim that (A, B_1, A', B'_1, I)

is an even ℓ -pretemplate partition of G_1 . Since $G_1[A \cup I \cup A']$ is exactly $G[A \cup I \cup A']$, conditions (3.), (4.) (5.), (6.) and (7.) hold. Adding the fact that $N_{G_1}(B_1) \subseteq A^* \cap V(G_1) = A$ by condition (b) of preblowup, condition (1.) for a pretemplate holds and symmetrically also condition (2.). Now condition (8.) holds because w^* and w'^* are universal in respectively $G^*[A \cup B_1]$ and $G^*[A' \cup B'_1]$. By (10), the last two conditions for a pretemplate are fulfilled by (A, B_1, A', B'_1, I) . Hence, by Lemma 8.8, G_1 is a an even ℓ -template. It is twinless by Lemma 8.9. We also notice that by construction w^* (respectively w'^*) belongs to G_1 and is universal in $G_1[A \cup B_1]$ (respectively $G_1[A' \cup B'_1]$). Since (A, B, A', B', I, w, w') is a proper ℓ -partition of G, there exist isolated vertices in $G^*[A]$ and $G^*[A']$. Hence $(A, B_1, A', B'_1, I, w^*, w'^*)$ is a proper ℓ -partition of G_1 .

We now prove that G^* is a proper blowup of G_1 .

By the definition of a preblowup and by (11), for all $u \in V(G_1)$, K_u is a clique and $V(G^*) = \bigcup_{u \in V(G_1)} K_u$

(12) If $u, v \in V(G_1)$ and $uv \notin E(G_1)$, then K_u is anticomplete to K_v .

If $u \in I$ or $v \in I$, the conclusion follows directly from condition (i) of preblowup. So we may assume up to symmetry that $u \in A \cup B_1$. By conditions (a) and (b) of preblowup, we may assume $v \in A \cup B_1$. If $u, v \in A$, then the result follows from (3), so we may assume that $v \in B_1$.

Now suppose for a contradiction that there exist $u^* \in K_u$ and $v^* \in K_v$ such that $u^*v^* \in E(G_1)$. By the choice of vertices in B_1 , for all $v^* \in K_v$, $N[v^*] \subseteq N[v]$. So $u^*v \in E(G_1)$. For the same reason or by (8), for all $u^* \in K_u$, $N[u^*] \subseteq N[u]$. Hence $uv \in E(G_1)$, a contradiction. This proves (12).

(13) If uv is a solid edge of G_1 then K_u is complete to K_v .

Otherwise, let $u^* \in K_u$ and $v^* \in K_v$ such that $u^*v^* \notin E(G)$. Since uv is a solid edge, up to symmetry, $u, v \in A$ or $u, v \in B_1$ or $u \in A$, $v \in B_1$ and in this last case u is not an isolated vertex of $G[H_v]$.

By (4) the case where u and v are in A cannot happen. Assume then that $v \in B_1$. By Lemma 8.3, there exist $a, b \in H_v$ (and hence in A) that are not adjacent. Assume that u is also in B_1 . Since u and v are adjacent, by (9) we may assume without loss of generality that $H_v \subseteq H_u$ and so aand b belong to H_u too. Then, by the definition of K_u and K_v , we get a C_4 induced by $\{u^*, v^*, a, b\}$, a contradiction.

So u should be in A, and to avoid a C_4 induced by $\{u^*, v^*, a, b\}$, u^* should be non-adjacent to at least one of a and b, say a. In particular, $a \neq u$. Then, by (4), $ua \notin E(G_1)$. So u does not belong to $N(H_v)$ and since uv is an edge of G_1 , we get that $u \in H_v$. Since uv is solid, u has at

least one neighbor in H_v , and we know that u is not adjacent to at least one vertex in H_v (namely a). Hence, as H_v is anticonnected, there exist non-adjacent vertices $c, d \in H_v$ such that $uc \notin E(G_1)$ and $ud \in E(G_1)$. Now $u^*P_{u^*}u'^*c'P_ccv^*du^*$ or $u^*P_{u^*}u'^*w'^*c'P_ccv^*du^*$ is a hole of length $2\ell + 1$, a contradiction again.

This proves (13).

(14) For all $u \in V(G_1)$ and $1 \le i \le j \le k_u$, $N[u_i] \subseteq N[u_j]$.

The result follows from how vertices are ordered after the proof of (6) (vertices in I), (8) (vertices in A or A') and (11) (vertices in B_1 or B'_1) and from (13). This proves (14).

(15) If uv is a flat edge of G_1 , then u is complete to K_v and v is complete to K_u .

By definition of a flat edge, either u and v are in I or one is in I and the other is in A or in A'. The result follows from (6), (8), and conditions (a2) (applied to A or A') and (i1) of the preblowup. This proves (15).

(16) If ux is an optional edge of G_1 with $u \in A$ and $x \in B_1$ (resp. $u \in A'$ and $x \in B'_1$), then u is complete to K_x .

The result follows from the definition of K_x when $x \in B_1$. This proves (16).

(17) If ux and uy are optional edges with $u \in A$, $x, y \in B_1$ and $H_y \subsetneq H_x$ (resp. $u \in A'$, $x, y \in B'_1$ and $H'_y \subsetneq H'_x$), then every vertex of K_u with a neighbor in K_y is complete to K_x .

Otherwise, let u^* be a vertex in K_u that has a neighbor y^* in K_y and a non-neighbor x^* in K_x . Since H_x and H_y are not disjoint, xy is a solid edge of G_1 and by (13), $x^*y^* \in E(G_1)$.

Since x and y are not equivalent, there exists a vertex a such that $a \in N_A(y) \setminus N_A(x)$ or $a \in N_A(x) \setminus N_A(y)$. In the first case, by definition of a template, $a \in A \setminus N_A[H_x]$. Then since $H_y \subsetneq H_x$ and H_x is a module of A we get that a is anticomplete to H_x and hence to H_y . So $a \notin N_A(y)$, a contradiction; we may then conclude that $a \in N_A(x) \setminus N_A(y)$

By definition of the cliques in B, $x^*a \in E(G^*)$ and $y^*a \notin E(G^*)$. Therefore, to avoid a C_4 induced by $\{x^*, y^*, u^*, a\}$, it should be that $u^*a \notin E(G^*)$.

Now $aP_aa'u'^*P_{u^*}u^*y^*x^*a$ or $aP_aa'w'^*u'^*P_{u^*}u^*y^*x^*a$ is a hole of length $2\ell + 1$ a contradiction. This proves (17).

(18) w^* (resp. w'^*) is a universal vertex of $G^*[\bigcup_{u \in A \cup B_1} K_u]$ (resp. $G^*[\bigcup_{u \in A' \cup B'_1} K_u]$).

By (1), w^* is complete to A^* and so to $\bigcup_{u \in A} K_u$. Furthermore, from the definition of G_1 we know that w^* is complete to $B_1 \setminus \{w^*\}$. Since all edges between vertices in B_1 are solid, by (13), w^* is complete to $B^* \setminus \{w^*\}$. Hence w^* is a universal vertex of $G^*[\bigcup_{u \in A \cup B} K_u]$. The proof for w'^* is symmetric. This proves (18).

From all the claims above, G^* satisfies all conditions to be a proper blowup of G_1 .

9 Graphs in $C_{2\ell}$ that contain a theta or a prism

The goal of this section is to prove the following.

Lemma 9.1 Let $\ell \geq 4$ be an integer. If G is a graph in $C_{2\ell}$ and G contains a theta or a prism, then one of the following holds:

- (a) G is a proper blowup of a twinless even ℓ -template;
- (b) G has a universal vertex;
- (c) G has a clique cutset.

The rest of this section is devoted to the proof of Lemma 9.1. So from here on, $\ell \geq 4$ is an integer and G is graph in $C_{2\ell}$ that contains a theta Θ or a prism Σ . By Lemma 3.10, the three paths of Θ have length ℓ and those of Σ have length $\ell - 1$. By Lemma 8.2, Θ and Σ are even ℓ -templates. Hence, we may define an integer k and a sequence F_0, F_1, F_2 of induced subgraphs of G as follows.

- k is the maximum integer such that G contains an even l-template with k principal paths. Observe that by Lemma 8.9, G in fact contains a twinless even l-template with k principal paths, because twins can be eliminated from templates by deleting hyperedges with equal vertexset while there are some.
- In G, pick a proper blowup F_1 of a twinless even ℓ -template F_0 with k principal paths. Note that F_0 exists and the proper ℓ -partition needed for the proper blowup exists by Lemma 8.11.
- Suppose that F_0 and F_1 are chosen subject to the maximality of the vertex-set of F_1 (in the sense of inclusion). Note that possibly F_0 is not a maximal template in the sense of inclusion, it can be that a smaller template leads to a bigger blowup (but F_0 has k principal paths).

• F_2 is obtained from F_1 by adding all vertices of $G \setminus F_1$ that are complete to F_1 .

Lemma 9.2 $V(F_2) \setminus V(F_1)$ is a (possibly empty) clique that is complete to F_1 .

Proof. Otherwise, G contains a C_4 .

We now introduce some notation. We denote by (A, B, A', B', I, w, w') the twinless proper even ℓ -partition that is used to blow up F_0 . We have $A = A_K \cup A_S$ and $A' = A'_K \cup A'_S$. When u is a vertex of F_0 , we denote by K_u the clique of F_1 that is blown up from u. We set $A^*_K = \bigcup_{u \in A_K} K_u$. We use a similar notation A^*_S , A'_K , A'_S , B^* , B'^* , I^* , A^* and A'^* .

9.1 Technical lemmas

We now prove lemmas that sum up several structural properties of G.

Lemma 9.3 If $u \in A \cup A' \cup I \cup \{w, w'\}$ and $v \in N_{V(F_0)}(u)$, then u is complete to K_v .

Proof. We prove this lemma using the conditions from the definition of blowups. If $u \in \{w, w'\}$, then the result follows from condition (h). If $u \in A \cup A'$, then the conclusion follows from conditions (d), (e) and (f). If $u \in I$, then the conclusion follows from condition (e).

Very often, Lemma 9.3 will be used in the following way. Suppose there exists a principal path $P = u \dots u'$ of F_0 . Suppose there exists a vertex x of P and $x^* \in K_x$. Then by Lemma 9.3 and condition (c) of blowups, $\{x^*\} \cup (V(P) \setminus \{x\})$ induces a path of F_1 . If $y \neq x$ is a vertex of P and $y^* \in K_y$, then $\{x^*, y^*\} \cup (V(P) \setminus \{x, y\})$ might fail to induce a path of F_1 , because it is possible that $xy \in E(G)$ while $x^*y^* \notin E(G)$. But under the assumption that $x^*y^* \in E(G)$ or $xy \notin E(G)$, we do have that $\{x^*, y^*\} \cup (V(P) \setminus \{x, y\})$ induces a path of F_1 . Several variant of this situation will appear soon and we will simply justify them by referring to Lemma 9.3.

When u is a vertex in A, we denote by P_u the unique principal path of F_0 that contains u. Its end in A' is then denoted by u'. We denote by u^+ the neighbor of u in P_u . We denote by u^{++} the neighbor of u^+ in $P_u \setminus u$. Note that $u^+ \in I$ and $u^{++} \in I \cup A'$ ($u^{++} \in A'$ if and only if $\ell = 4$ and $u \in A_S$).

For any distinct $u, v \in A$, from the definition of even templates, exactly one of $V(P_u) \cup V(P_v)$ or $V(P_u) \cup V(P_v) \cup \{w\}$ or $V(P_u) \cup V(P_v) \cup \{w'\}$ or $V(P_u) \cup V(P_v) \cup \{w, w'\}$ induces a hole that is denoted by $C_{u,v}$. Such a hole is called a *principal hole*.

So, there are three kinds of principal holes: those that contain exactly one of w and w', those that contain none of w and w' and those that contain both w and w'. Recall that by Lemma 8.7, some holes of a template contain two principal paths plus one or two extra vertices, but it may fail to be a principal hole (because the extra vertices may fail to be w or w'). Though we do not use this information formally, it is worth noting that by Lemma 9.3, when C is a principal hole, $\bigcup_{v \in V(C)} K_v$ induces a ring. But when C is a non-principal hole, it may happen that $\bigcup_{v \in V(C)} K_v$ does not induce a ring (because there might be in C an optional edge uv with $u \in A$ and $v \in B$, and after the blowup process, there might be that no vertex in K_v is complete to K_u).

Lemma 9.4 If $u \in V(F_0)$ and $u^* \in K_u$, then u^* has two neighbors in $V(F_0) \setminus K_u$ that are not adjacent.

Proof. If $u \in I$, then let P be the principal path that contains u. By Lemma 9.3, u^* is adjacent to the two neighbors of u in P.

If $u \in A \cup A'$, say $u \in A$ up to symmetry, then we claim that u has a neighbor z in $A \cup B$. This is clear if u is not isolated in A and otherwise we set z = w. By Lemma 9.3, z and u^+ are non-adjacent neighbors of u^* .

If $u \in B$, then by the definition of a template, H_u contains two non adjacent vertices a and b that are neighbors of u. By Lemma 9.3, a and b are both adjacent to u^* .

Lemma 9.5 If K is a clique of F_0 , $K^* = \bigcup_{v \in K} K_v$ and D is a connected induced subgraph of $G \setminus F_2$ such that $N_{V(F_1)}(D) \subseteq K^*$, then $N_{V(F_1)}(D)$ is a clique.

Proof. For suppose not. This means that there exists $u^*, v^* \in K^*$ and $x_u, x_v \in D$ such that $u^*v^* \notin E(G)$ and $x_uu^*, x_vv^* \in E(G)$ (possibly $x_u = x_v$). Since D is connected, there exists a path P in D from x_u to x_v . Suppose that u^*, x_u, v^*, x_v and P are chosen subject to the minimality of P. It follows that $u^*x_uPx_vv^*$ is a path, and recall that by assumption its interior is anticomplete to $F_1 \setminus K^*$.

Since $u^*v^* \notin E(G)$, u^* and v^* are in different blown-up cliques. Denote by K_u and K_v the blown-up cliques such that $u^* \in K_u$ and $v^* \in K_v$. By hypothesis, $u, v \in K$ and so $uv \in E(G)$. Since $u^*v^* \notin E(G)$, by condition (d) of blowups, uv is not a solid edge of G. If uv is a flat edge of F_0 , then uv is an edge of a principal path and we know that this principal path belongs to a principal hole C. Note that apart from u and v, no vertex of C is in K since K is a clique. By Lemma 9.3, in G, $(\{u^*, v^*\}) \cup V(C)) \setminus \{u, v\}$ induces a path Q of length $2\ell - 1$. So P and Q form a hole of length at least $2\ell + 1$, a contradiction.

If uv is an optional edge of F_0 , say with $u \in A$ and $v \in B$, then $u \in H_v$, and there exists a in H_v such that $au \notin E(F_0)$. Therefore, P_u , P_a , v and possibly w' form a hole C^* . By condition (f) of blowups (if va is optional), or by condition (d) (if va is solid), a is complete to K_v . By Lemma 9.3 it follows that $(\{u^*, v^*\}) \cup V(C^*)) \setminus \{u, v\}$ induces a path Q of length $2\ell - 1$. So P and Q form a hole of length at least $2\ell + 1$, a contradiction again. \Box

Lemma 9.6 If $x \in V(G) \setminus V(F_2)$ and C is a principal hole of F_0 , then x is minor w.r.t. C.

(We remind that x is minor w.r.t. C if the neighborhood of x in C is included in a 3-vertex path of C.)

Proof. Otherwise let $C = C_{u,v}$ for some $u, v \in A$ such that x is major w.r.t. C. By Lemma 3.10, x and C form a universal wheel.

(1) x is complete to all principal paths.

We know already that x is complete to P_u and P_v . Let $P_t = t \dots t'$ be a principal path where $t \neq u, v$. If one of u, v, say u, is in A_K then P_u contains at least 4 vertices, so x is major w.r.t. $C_{u,t}$ and as above we may conclude that x and $C_{u,t}$ form a universal wheel. Consider now the case where both u and v are in A_S . Then C contains w and w' and $C_{u,t}$ contains at least one of w, w'. So x is adjacent to at least 4 vertices of $C_{u,t}$ and we may again conclude that x is complete to P_t . This proves (1).

(2) x is complete to $B \cup B'$.

Let $y \in B \cup B'$. By definition of a template, y has two neighbors a and b, both in A or both in A', that are non-adjacent. Therefore a, b, y and x form a C_4 , unless x is adjacent to y. This proves (2).

By (1) and (2), x is complete to $I \cup A \cup A' \cup B \cup B' = V(F_0)$.

Let z be a vertex of F_0 and $z^* \in K_z$. By Lemma 9.4, there exists $a, b \in V(F_0)$ such that $z^*a, z^*b \in E(G)$ and $ab \notin E(G)$. Since there is no C_4 in G it should be that $xz^* \in E(G)$. This proves that x is complete to F_1 . Hence, $x \in V(F_2)$, a contradiction.

Lemma 9.7 Let a and b be two non-adjacent vertices of some principal hole C of F_0 . If some vertex x of $V(G) \setminus V(F_2)$ has neighbors in both K_a and K_b , then a and b have a common neighbor c in C, x is adjacent to c, and x is anticomplete to every K_d such that $d \in V(C) \setminus \{a, b, c\}$.

Proof. Let $a^* \in K_a$ and $b^* \in K_b$ be two neighbors of x. Since $ab \notin E(G)$, by Lemma 9.3, $\{a^*, b^*\} \cup V(C) \setminus \{a, b\}$ induces a hole C^* . Since x is adjacent to a^* and b^* , by Lemma 3.10, x has another neighbor c in C^* (and in fact in C since $c \neq a^*, b^*$). If c is not adjacent to a^* or b^* , then x is major w.r.t. C^* , so by Lemma 3.10, C^* and x form a universal wheel. It follows that xis major w.r.t. C, a contradiction to Lemma 9.6.

We proved that a and b have a common neighbor c in C and that x is adjacent to c. Suppose for a contradiction that x has a neighbor $d^* \in K_d$ where $d \in V(C) \setminus \{a, b, c\}$. By the same argument as above, since x has neighbors in K_d and K_c , c and d must have a common neighbor in C, and this common neighbor must be a or b, say a up to symmetry. So, x has neighbors in K_d and K_b while b and d have no common neighbors in C, so we may reach a contradiction as above.

9.2 Connecting vertices of a template

Lemma 9.8 If $x \in A \cup B$ and $y \in A' \cup B'$, then there exists in G a path P of length $\ell - 2$, $\ell - 1$, ℓ or $\ell + 1$ from x to y that contains a principal path. More specifically:

- If $x \in A$ and $y \in A'$, then P has length $\ell 2$, $\ell 1$, ℓ or $\ell + 1$.
- If $x \in A$ and $y \in B'$, or if $x \in B$ and $y \in A'$, then P has length $\ell 1$, ℓ or $\ell + 1$.
- If $x \in B$ and $y \in B'$, then P has length ℓ or $\ell + 1$. Furthermore in that case, there exists another path Q from x to y of length ℓ or $\ell + 1$ containing a principal path and Q contains no interior vertex of P.

Proof. Suppose first that $x \in A$, say $x = v_i$. If $y \in A'$, then set $y = v'_j$. If i = j, then P_i has length $\ell - 2$ or $\ell - 1$. Assume $i \neq j$. If $v_i v_j$ or $v'_i v'_j$, say $v_i v_j$ is an edge then $v_i v_j P_{v_j} v'_j$ is a path of length $\ell - 1$ or ℓ . If none of $v_i v_j$ or $v'_i v'_j$ is an edge then v_i and v_j are in A_S and $v_i P_{v_i} v'_i w' v'_j$ is a path of length ℓ . If $y \in B'$, then one of $v_i P_i v'_i y$ or $v_i P_i v'_i w' y$ is the path of length at most $\ell + 1$ we are looking for. The proof is similar when $y \in A'$.

We may therefore assume that $x \in B$ and $y \in B'$. Assume first that there exist v_i and v_j in A such that x is adjacent to v_i and v_j and y is adjacent

to v'_i and v'_j . Then $xv_iP_{v_i}v'_iy$ and $xv_jP_{v_j}v'_jy$ are two paths of length ℓ or $\ell + 1$. Consider now the case when there exists one vertex v_i in H_x such that v'_i belongs to H_y . Since both $G[H_x]$ and $G[H_y]$ are anticonnected and contain at least two vertices, there exists $u \in H_x$ non adjacent to v_i and $v' \in H_y$ non adjacent to v'_i . We may assume that u' is not adjacent to y and that v is not adjacent to x since else we are in the previous case. So now v should be anticomplete to H_x and u' should be anticomplete to H_y . So $\{v_i, u, v\}$ and $\{v'_i, u', v'\}$ are both stable sets of G, this is possible if and only if $\{v_i, u, v\} \subseteq A_S$. We have then two paths $xuP_uu'w'y$ and $xwvP_vv'y$ of length $\ell + 1$. It remains to consider the case where each $v_i \in H_x$ is such that $v'_i \notin H_y$. By definition each of H_x and H_y contains a pair of non adjacent vertices and hence there exist two distinct vertices u, v' such that $u \in H_x \cap A_S$ and $v' \in H_y \cap A'_S$. We have then again two paths $xuP_uu'w'y$ and $xwvP_uv'y$ of length $\ell + 1$.

9.3 Connecting vertices of F_1

We here explain how lemmas of Subsection 9.2 are extended from F_0 to F_1 .

Lemma 9.9 If $u^* \in A^* \cup B^*$ and $v^* \in A'^* \cup B'^*$, then there exists in F_1 a path P^* of length $\ell - 2$, $\ell - 1$, ℓ or $\ell + 1$ from u^* to v^* that contains the interior of a principal path P. More specifically:

- If $u^* \in A^*$ and $v^* \in A'^*$, then P^* has length $\ell 2$, $\ell 1$, ℓ or $\ell + 1$.
- If $u^* \in A^*$ and $v^* \in B'^*$, or if $u^* \in B^*$ and $v^* \in A'^*$, then P^* has length $\ell 1$, ℓ or $\ell + 1$.
- If u^{*} ∈ B^{*} and v^{*} ∈ B'^{*}, then P^{*} has length l or l + 1. Furthermore in that case there exists another path Q^{*} from u^{*} to v^{*} of length l or l + 1 which contains the interior of a principal path Q ≠ P.

Proof. Let u and v be such that $u^* \in K_u$ and $v^* \in K_v$. Let P be a path in F_0 like in Lemma 9.8 from u to v (so P contains the interior of some principal path Q). By Lemma 9.3, $\{u^*, v^*\} \cup V(P) \setminus \{u, v\}$ induces a path of the same length as P that contains the interior of Q. \Box

Lemma 9.10 If in G some vertex x is adjacent to the ends of a path P of length at most $\ell + 1$ not containing x, then x is complete to V(P).

Proof. Otherwise, a shortest cycle in $G[V(P) \cup \{x\}]$ has length at least 4 and at most $\ell + 3$. Since $\ell \ge 4$ implies $\ell + 3 < 2\ell$, this is a contradiction. \Box

9.4 Attaching a vertex to F_1

In this subsection, we show that for all vertices x of $V(G) \setminus V(F_2)$, $N_{V(F_1)}(x)$ is a clique (see Lemma 9.14).

Lemma 9.11 If $x \in V(G) \setminus V(F_2)$ has no neighbor in I^* , then $N_{V(F_1)}(x)$ is a clique.

Proof. Suppose for a contradiction that $N_{V(F_1)}(x)$ is not a clique.

(1) We may assume that $N_{V(F_1)}(x) \subseteq A^* \cup B^*$.

If x has neighbors in both $A^* \cup B^*$ and $A'^* \cup B'^*$, then consider a path P as in Lemma 9.9 from a neighbor of x in $A^* \cup B^*$ to a neighbor of x in $A'^* \cup B'^*$. By Lemma 9.10, x is complete to V(P). This is a contradiction since x has no neighbor in I^* . Hence x does not have neighbors in both $A^* \cup B^*$ and $A'^* \cup B'^*$, and our claim follows up to symmetry. This proves (1).

(2) There exist non-adjacent $a, b \in A$ such that x has neighbors in both K_a and K_b .

By Lemma 9.5, since $N_{V(F_1)}(x)$ is not a clique, there should exist two non-adjacent vertices $a, b \in V(F_0)$ such that x has a neighbor $a^* \in K_a$ and a neighbor $b^* \in K_b$. By (1), $a, b \in A \cup B$.

If $a, b \in A$, then our conclusion holds, so we may assume that $b \in B$.

If $a \in A$, then since $ab \notin E(G)$, H_b is anticomplete to a. Let P_a^* be the path induced by $\{a^*\} \cup (V(P_a) \setminus \{a\})$. Let $v \in H_b$. We may assume that $xv \notin E(G)$ for otherwise our claim holds (with a and v). Note that since $ab, av \notin E(G)$, by (c) of blowup, $a^*b^*, a^*v \notin E(G)$. Now, $a^*xb^*vP_vv'a'P_a^*a^*$ (in case one of a, v belongs to A_K) or $a^*xb^*vP_vv'a'P_a^*a^*$ (in case both a, v belong to A_S) form a hole of length $2\ell + 1$, a contradiction. Hence, we may assume $a \in B$.

Since $ab \notin E(G)$, by Lemma 8.4, $\{a\} \cup H_a$ is anticomplete to $\{b\} \cup H_b$. We may assume that x is anticomplete to $H_a \cup H_b$ for otherwise we may apply the proofs above. Hence, for $u \in H_a$ and $v \in H_b$, $ua^*xb^*vP_vv'u'P_uu$ or $ua^*xb^*vP_vv'w'u'P_uu$ is a hole of length $2\ell + 2$. This proves (2).

Now the sets K_u for all $u \in A \cup A' \cup I$, $B^* \cup \{x\}$ and B'^* form a preblowup of F_0 . All conditions are easily checked. In particular x satisfies condition (b) by (1) and (b2) by (2)). So, by Lemma 8.15, $G[V(F_1) \cup \{x\}]$ is

a proper blowup of some ℓ -template with k principal paths. This contradicts the maximality of F_1 .

Lemma 9.12 If there exist $x \in V(G) \setminus V(F_2)$ and $u \in A$ such that x has neighbors in both K_u and K_{u^+} and is anticomplete to $K_{u^{++}}$, then $N_{V(F_1)}(x)$ is a clique.

Proof. Suppose for a contradiction that $N_{V(F_1)}(x)$ is not a clique.

(1) x is anticomplete to $A'^* \cup B'^* \cup (I^* \setminus K_{u^+})$.

If x has a neighbor t^* in some K_t such that $t \in (A' \cup I) \setminus \{u^+\}$, then note that $t \neq u^{++}$ by assumption. Let C be a principal hole that contains t and u. By Lemma 9.7 applied to t and u (that are by definition non adjacent in C) we should have that u^+ is a common neighbor of u and t in C. This is not possible since the neighbors of u^+ in C are u and $u^{++} \neq t$. So x is anticomplete to $A'^* \cup (I^* \setminus K_{u^+})$.

It remains to prove that x is anticomplete to B'^* . Otherwise, x has a neighbor $t \in B'^*$. Consider a path P from t to a neighbor of x in K_u as in Lemma 9.9 and let Q be the principal path whose interior is contained in P. By Lemma 9.10, x is complete to V(P) and hence to V(Q). This is impossible since we have shown that x is anticomplete to A'^* . This proves (1).

From here on, u^* and u^{+*} are neighbors of x in respectively K_u and K_{u^+} .

(2) x has a neighbor $y^* \in B^*$ that is complete to A^* .

Note that x has a neighbor $y^* \in K_y$ for some $y \in A \cup B \setminus \{u\}$, for otherwise, by (1), $N_{V(F_1)}(x) \subseteq K_u \cup K_{u^+}$ and by Lemma 9.5, $N_{V(F_1)}(x)$ is a clique, a contradiction. In case $y^* = w$ we are done, so from now on we will assume that w is not adjacent to x and $y^* \neq w$.

Let $v \in A \setminus \{u\}$ and assume that there exists $v^* \in K_v$ which not adjacent to y^* . Then $xv^* \notin E(G)$ for otherwise $\{x, y^*, w, v^*\}$ induces a C_4 . Let P_v^* be the path induced by $\{v^*\} \cup (V(P_v) \setminus \{v\})$. Either $xy^*wv^*P_v^*v'w'u'P_uu^{++}u^{+*}x$ (in case $v'u' \notin E(G)$) or $xy^*wv^*P_v^*v'u'P_uu^{++}u^{+*}x$ (in case $v'u' \in E(G)$) is a hole of length at least $2\ell + 1$. In both cases we get a contradiction. So we have shown that for every $v \in A \setminus \{u\}, y^*$ is complete to K_v and hence y is complete to $A \setminus \{u, y\}$.

We also have that $u^*y^* \in E(G)$ for otherwise, $\{x, y^*, w, u^*\}$ induces a C_4 . So, if $y^* \in A^*$ then A contains no isolated vertex, a contradiction to Lemma 8.10. So $y^* \in B^*$. If y^* has a non-neighbor $v^* \in A^*$ then from what precedes we have $v^* \in K_u$. Now there exists a path Q of length 1, 2 or 3 from x to v^* with interior in K_{u^+} (either xv^* or $xu^{+*}v^*$, or xu^+v^* or

 $xu^{+*}u^+v^*$). Hence, xQv^*wy^*x is a hole of length 4, 5 or 6, a contradiction. This proves (2).

(3) $N_A(x) \setminus \{u\} = N_A(u).$

If there exists $v \in N_A(x) \setminus N_A[u]$, then $vP_v v'u'P_u u^{++}u^{+*}xv$ (in case exactly one of u, v is in A_K) or $vP_v v'w'u'P_u u^{++}u^{+*}xv$ (in case u, v are both in A_S) is a hole of length $2\ell - 1$, a contradiction.

Conversely, suppose there exists $v \in N_A(u) \setminus N_A(x)$.

Then $v'P_vvy^*xu^{+*}u^{++}P_uu'w'v'$ (in case exactly one of u, v is in A_K) or $v'P_vvy^*xu^{+*}u^{++}P_uu'v'$ (in case u, v are both in A_K) is a hole of length $2\ell + 1$, a contradiction. This proves (3).

(4) x is complete to K_u .

Suppose there exists $r \in K_u$ such that $rx \notin E(G)$.

If $xu^+ \in E(G)$ then $\{r, y^*, u^+, x\}$ induces a C_4 , a contradiction. Hence $xu^+ \notin E(G)$. Now by condition (e) of blowups, either $\{x, y^*, r, u^{+*}\}$ induces a C_4 or $\{x, y^*, r, u^+, u^{+*}\}$ induces a C_5 . This proves (4).

Now, the sets K_v for all $v \in (A \setminus u) \cup I \cup A'$, $K_u \cup \{x\}$, B^* and B'^* form a preblowup of F_0 . All conditions are easy to check. In particular, $K_u \cup \{x\}$ is a clique by (4), conditions (a), (b) and (i) follows from (1), condition (a1) from (3), condition (b1) from (2) and condition (a2) from our assumptions.

Hence by Lemma 8.15, $G[V(F_1) \cup \{x\}]$ is a proper blowup of some twinless odd ℓ -template with k principal paths that is an induced subgraph of G a contradiction to the maximality of F_1 .

Lemma 9.13 If $x \in V(G) \setminus V(F_2)$ has no neighbor in $B^* \cup B'^*$, then $N_{V(F_1)}(x)$ is a clique.

Proof. Suppose for a contradiction that $N_{V(F_1)}(x)$ is not a clique. By Lemma 9.11, x has a neighbor in I^* . So x has a neighbor in a clique blown up from an internal vertex of some principal path $P_v = v \dots v'$. Let a (resp. b) be the vertex of P_v closest to v (resp. to v') along P_v and such that x has a neighbor in K_a (resp. K_b).

Suppose first that a = b, so $a \in I$ and a is the only vertex of P_v whose clique contains a neighbor of x. Hence, as $N_{V(F_1)}(x)$ is not a clique, x has a neighbour in some K_y with $y \in V(F_0) \setminus V(P_v)$, and since by assumption x has no neighbor in $B^* \cup B'^*$, $y \in A \cup A' \cup I$. So, y and a are nonadjacent members of some principal hole. By Lemma 9.7, x has a neighbor in some clique K_c where c is adjacent to both a and y, a contradiction to the properties implied by the supposition that a = b.

Suppose now that $ab \in E(G)$. Then a and b are the only vertices of P_v whose cliques contain a neighbor of x. If both a and b are internal vertices of P_v , then as in the previous paragraph, we can show that no neighbour of x is in some K_y with $y \in V(F_0) \setminus V(P_v)$ and hence $N_{V(F_1)}(x) \subseteq K_a \cup K_b$. So, by Lemma 9.5, $N_{V(F_1)}(x)$ is a clique, a contradiction. It follows that at least one of a or b is an end of P_v . Up to symmetry, we may assume that a = v and $b = v^+$. Note that, by the definition of b, x is then anticomplete to $K_{v^{++}}$. Hence, by Lemma 9.12, $N_{V(F_1)}(x)$ is a clique, a contradiction.

Hence, $a \neq b$ and $ab \notin E(G)$. So, by Lemma 9.7, a and b have a common neighbor u in P_v . So, a, u and b are consecutive along P_v (in particular, $u \in I$).

(1) x is complete to K_u .

Otherwise, let $u^* \in K_u$ be non-adjacent to x. There exists a path Q_a of length 2 or 3 from u^* to x with interior in K_a (either xa^*u^* , or xa^*au^* for some a^* in K_a). There exists a similar path Q_b . So, Q_a and Q_b form a hole of length 4, 5 or 6, a contradiction. This proves (1).

(2) x is anticomplete to $V(F_1) \setminus (K_a \cup K_u \cup K_b)$.

This follows from Lemma 9.7 and from the assumption that x is anticomplete to $B^* \cup B'^*$. This proves (2).

(3) x has neighbors in each of K_a , K_b .

This follows from the definition of a and b. This proves (3).

Now the sets K_v for all $v \in (A \cup A' \cup I) \setminus \{u\}$, $K_u \cup \{x\}$, B^* and B'^* form a preblowup of F_0 . All conditions are easily checked, in particular $K_u \cup \{x\}$ is a clique by (1), it satisfies condition (i) by (2) and condition (i1) by (3).

Hence, by Lemma 8.15 $G[V(F_1) \cup \{x\}]$ is a proper blowup of some twinless odd ℓ -template with k principal paths that is an induced subgraph of G. This contradicts the maximality of F_1 .

Lemma 9.14 For all vertices x of $V(G) \setminus V(F_2)$, $N_{V(F_1)}(x)$ is a clique.

Proof. Suppose for a contradiction that $N_{V(F_1)}(x)$ is not a clique.

(1) There exists a principal path $P_u = u \dots u'$ of F_0 such that x is anticomplete to $I^* \setminus \bigcup_{v \in V(P_u)} K_v$.

Otherwise, there exist two distinct principal paths P and Q of F_0 such that a in the interior of P, b in the interior of Q and x has neighbors in both K_a and K_b . Note that then a and b are non adjacent and do not share any neighbour. This contradicts Lemma 9.7, applied to the principal hole C of F_0 containing P and Q. This proves (1).

(2) We may assume that x has no neighbor in $B^{\prime*}$ and has a neighbor $y^* \in K_y$ where $y \in B$.

Suppose that x has a neighbor $u^* \in B^*$ and a neighbor $v^* \in B'^*$. Let P and Q be as in Lemma 9.9. By Lemma 9.10, x is complete to both V(P) and V(Q). In particular, x has neighbors in the interior of two distinct principal paths, a contradiction to (1). So, up to symmetry, we may assume that x has no neighbor in B'^* . Hence, by Lemma 9.13, x has neighbors in B^* . This proves (2).

(3) x is adjacent to u and u^+ and has a neighbor in $K_{u^{++}}$. Moreover, x is anticomplete to $(A^* \cup I^* \cup A'^* \cup B'^*) \setminus (K_u \cup K_{u^+} \cup K_{u^{++}})$.

By Lemma 9.11, x has at least one neighbor in I^* and by (1), such a neighbor is in a clique blown up from an internal vertex of P_u . So, let v be the vertex of P_u closest to u' along P_u such that x has a neighbor $v^* \in K_v$. So $v \neq u$ and $v \in A' \cup I$. We set $Q = y^* u P_u v$ if $y^* u \in E(G)$ and $Q = y^* w u P_u v$ otherwise. Let Q^* be the path induced by $\{v^*\} \cup (V(Q) \setminus \{v\})$ and observe that Q^* has length at most $\ell + 1$. By Lemma 9.10, x is complete to Q^* . If $v \notin \{u^+, u^{++}\}$, then x has neighbors in at least 4 cliques blown up from vertices of P_u and this contradicts Lemma 9.7. If $v = u^+$, x is adjacent to u (since x is complete to Q^*) and anticomplete to $K_{u^{++}}$, so by Lemma 9.12, $N_{V(F_1)}(x)$ is a clique, a contradiction. So, $v = u^{++}$, meaning that x is adjacent to u and u^+ , and is anticomplete to $I^* \setminus (K_{u^+} \cup K_{u^{++}})$ by (1).

If x has neighbors in some K_a for $a \in A \setminus \{u\}$ then x and $C_{u,a}$ contradict Lemma 9.7. Hence x is anticomplete to $A^* \setminus \{K_u\}$.

By (2), x is anticomplete to B'^* . It remains to check that x is anticomplete to $A'^* \setminus K_{u^{++}}$. So, suppose x has a neighbor z^* in some K_z where $z \in A' \setminus \{u^{++}\}$. Then a principal hole that contains z and u contradicts Lemma 9.7. This proves (3).

Let u^{++*} be a neighbor of x in $K_{u^{++}}$ and P_u^* be the path induced by $(V(P_u) \setminus \{u^{++}\}) \cup \{u^{++*}\}.$

(4) For every $z \in B$ such that x is adjacent to some z^* in K_z we have $N_A(z) = N_A[u]$ (in particular $N_A(y) = N_A[u]$).

Suppose there exists $v \in N_A(z) \setminus N_A[u]$. By condition (d) or (f) of blowups, $vz^* \in E(G)$. So, by (3), $xz^*vP_vv'u'P_u^*u^{++*}x$ (if one of $u, v \in A_K$) or $xz^*vP_vv'w'u'P_u^*u^{++*}x$ (if both u and v are in A_S) is a hole of length $2\ell - 1$, a contradiction. This proves that $N_A(z) \subseteq N_A[u]$. In particular, u has at least one neighbor in H_z , so by condition (k) of templates, $uz \in E(G)$.

Suppose there exists $v \in N_A(u) \setminus N_A(z)$ (so $z \neq w$). By condition (c) of blowups, $vz^* \notin E(G)$. By (3), $xv \notin E(G)$. Hence $xw \in E(G)$, for otherwise $xz^*wvP_vv'u'P_u^*u^{++*}x$ or $xz^*wvP_vv'u'P_u^*u^{++*}x$ is a hole of length $2\ell + 1$. Since the partition is proper there exists a vertex $c \in A_S$ which is isolated in G[A] and, since $v \in N(u)$, $c \neq u$. Again by (3), $xc \notin E(G)$ and $xwcP_cc'u'P_u^*u^{++*}x$ (in case $u \in A_K$) or $xwcP_cc'w'u'P_u^*u^{++*}x$ (in case $u \in A_S$) is a hole of length $2\ell - 1$, a contradiction. This proves (4).

(5)
$$N_{F_1}(x) \subseteq K_{u^{++}} \cup K_{u^+} \cup K_u \cup K_y$$

By (3) $N_{F_1}(x) \subseteq K_{u^{++}} \cup K_{u^+} \cup K_u \cup B^*$. Suppose there exists $z^* \in K_z$ such that $xz^* \in E(G)$ and $z \in B \setminus \{y\}$. By (4), $N_A(z) = N_A[u]$ and $N_A(y) = N_A[u]$. So, by Lemma 8.9, y and z are twins of F_0 , a contradiction. This proves (5).

(6) $y \neq w$.

If y = w, then by (4), $N_A(w) = N_A[u] = A$ and so u is a universal vertex of G[A]. By Lemma 8.10, there exists also at least one isolated vertex in G[A], a contradiction to $|A| \ge 3$. This proves (6).

(7) $N_{K_u}(x)$ is complete to $N_A[u]$.

By (4), $N_A(y) = N_A[u]$. The result follows from conditions (d) and (f) of blowups. This proves (7).

(8) x is complete to K_{u^+} .

By (3), $ux \in E(G)$. Suppose for a contradiction that there exists $u^{+*} \in K_{u^+}$ non-adjacent to x. By condition (e) of blowups, $u^{+*}u, u^{+*}u^{++} \in E(G)$. Hence $xu^{++} \notin E(G)$ for otherwise $\{x, u^{++}, u^{+*}, u\}$ induces a C_4 . But now, either $\{x, u^{++*}, u^{+*}, u\}$ induces a C_4 (if $u^{+*}u^{++*} \in E(G)$) or $\{x, u^{++*}, u^{+*}, u\}$ induces a C_5 (if $u^{+*}u^{++*} \notin E(G)$), a contradiction. This proves (8).

(9) $K_u \cup K_y$ is a clique.

Since by (4) $N_A(y) = N_A[u]$, *u* cannot be an isolated vertex of H_y . Hence, *uy* is a solid edge. So, by condition (d) of blowups, K_u is complete K_y . This

proves (9).

We define $B_0 = B^* \setminus N_{K_u}(x)$.

Now the sets K_v for all $v \in (A \cup I \cup A') \setminus \{u, u^+\}, K_u \cup N_{K_y}(x), K_{u^+} \cup \{x\}, B_0$ and B'^* form a preblowup of F_0 . All conditions are easy to check. In particular, $K_u \cup N_{K_y}(x)$ is a clique by (9), $K_{u^+} \cup \{x\}$ is a clique by (8), conditions (a), (b) and (i) follows from (5), condition (a1) from (7), condition (a2) holds because x is complete to $N_{K_y}(x)$, condition (i1) follows from (3) and condition (b1) holds because of (6).

Hence, by Lemma 8.15, $G[V(F_1) \cup \{x\}]$ is a proper blowup of some twinless odd ℓ -template with k principal paths that is an induced subgraph of G, a contradiction to the maximality of F_1 .

9.5 Attaching a component

Lemma 9.15 If D is a connected component of $G \setminus F_2$, then N(D) is a clique.

Proof. Suppose that N(D) is not a clique. Since D is a connected component of $G \setminus F_2$ we have that $N(D) = N_{V(F_1)}(D)$. By Lemma 9.2, $N_{V(F_1)}(D)$ is not a clique. So, there exist a and b in D such that $N_{V(F_1)}(a) \cup N_{V(F_1)}(b)$ is not a clique, and a path P from a to b in D. We choose a and b subject to the minimality of the length of P. By Lemma 9.14, $a \neq b$ (so P has length at least 1).

We set $S_a^* = N_{V(F_1)}(a)$ and $S_b^* = N_{V(F_1)}(b)$. By Lemma 9.14, S_a^* and S_b^* are both cliques. Note that possibly $S_a^* \cap S_b^* \neq \emptyset$. We denote by $\operatorname{int}(P)$ the set of the internal vertices of P. We set $S_o^* = N_{V(F_1)}(\operatorname{int}(P))$.

We set $S_a = \{t \in V(F_0) : S_a^* \cap K_t \neq \emptyset\}$. We define S_b and S_\circ similarly. Note that S_a is possibly not included in S_a^* , and the same remark holds for S_b and S_\circ .

(1) There exist non-adjacent $x_a^* \in S_a^*$ and $x_b^* \in S_b^*$. Moreover, for all such x_a^* and x_b^* , $x_a^* aPbx_b^*$ is a path.

The existence of x_a^* and x_b^* follows from the definition of a and b, and $x_a^* a P b x_b^*$ is a path because of the minimality of P. This proves (1).

(2) $S_a^* \cup S_o^*$ and $S_b^* \cup S_o^*$ are cliques (in particular, S_o^* is a (possibly empty) clique of F_1 that is complete to both $S_a^* \setminus S_o^*$ and $S_b^* \setminus S_o^*$).

If $S_a^* \cup S_o^*$ is not a clique, then let x^*y^* be a non-edge in $S_a^* \cup S_o^*$. Since S_a^* is a clique by Lemma 9.14, we may assume $y^* \in S_o^*$. By definition of S_o^* , y^*

has a neighbor in int(P), and then x^*, y^* and some subpath of P contradict the minimality of P. The proof is similar for $S_b^* \cup S_o^*$. This proves (2).

Note that while $S_a^* \cup S_b^*$ is not a clique by assumption, it might be that $S_a \cup S_b$ is a clique (for instance when $S_a = \{u\}$, $S_b = \{v\}$ and uv is an optional edge of F_0).

(3) If $S_{\circ}^* \neq \emptyset$ then any two non adjacent vertices $x \in S_a^*$ and $y \in S_b^*$ are at distance 2 in F_1 .

Let $s^* \in S^*_{\circ}$. By (2), $s^* \neq x, y$ and xs^*y is a path in F_1 . This proves (3).

(4) $S_a \cup S_\circ$ and $S_b \cup S_\circ$ are cliques of F_0 (in particular, S_a and S_b are (non-empty) cliques of F_0 and S_\circ is a (possibly empty) clique of F_0 that is complete to both $S_a \setminus S_\circ$ and $S_b \setminus S_\circ$).

If $S_a \cup S_\circ$ is not a clique, then let xy be a non-edge of $S_a \cup S_\circ$. Since $x \in S_a \cup S_\circ$, there exists $x^* \in K_x \cap (S_a^* \cup S_\circ^*)$ and $y^* \in K_y \cap (S_a^* \cup S_\circ^*)$. By condition (c) of blowups, since $xy \notin E(G)$, K_x is anticomplete to K_y . So, $x^*y^* \notin E(G)$, a contradiction to (2).

The proof is similar for $S_b \cup S_o$. This proves (4).

For the next claim we use the path P defined at the very beginning of the proof.

(5) If a hole C of F_1 contains two non adjacent vertices $x \in S_a^*$ and $y \in S_b^*$, then P and C form either:

- a theta, and $S_a^* \cap V(C) = \{x\}$, $S_b^* \cap V(C) = \{y\}$ and the three paths, all of length ℓ , are the two paths between x and y in C and the path between x and y obtained by adding the edges ax and by to P; or
- a prism, $S_a^* \cap V(C) = \{x, z\}$, $S_b^* \cap V(C) = \{y, t\}$, the triangles of the prism are axz and byt and the three disjoint paths all of length $\ell 1$, are either:
 - the path P, the shortest path between x and y in C and the shortest path between z and t in C; or
 - the path P, the shortest path between x and t in C and the shortest path between z and y in C.

Note that since S_a^* is a clique, $S_a^* \cap V(C)$ contains x and at most one other vertex which should be adjacent to x. The same holds for S_b^* and y.

Let us assume that $S^*_{\circ} \cap V(C) \neq \emptyset$. Then by (3), there exists a unique vertex $t \in S^*_{\circ} \cap V(C)$, and t is such that $S^*_a \cap V(C) \subseteq \{x, t\}$ and $S^*_b \cap V(C) \subseteq$

 $\{y, t\}$. Hence C and P form a proper wheel centered at t, a contradiction to Lemma 3.10. So, $S_{\circ}^* \cap V(C) = \emptyset$.

If a and b have a common neighbor t in C, then x and y are the two neighbors of t in C and so, C and P form a proper wheel centered at t, again a contradiction to Lemma 3.10. So the neighborhoods of a and b in C are disjoint.

From the remarks above and Lemma 3.10, we obtain that C and P form a theta whose three paths have length ℓ or a prism whose three paths have length $\ell - 1$. This can happen only if we are in one of the three cases described in (5).

This proves (5).

(6) $S_a \cap I = S_b \cap I = \emptyset$.

Otherwise, up to symmetry, $S_a \cap I \neq \emptyset$. So, there exists a principal path $P_u = u \dots u'$ of F_0 whose interior intersects S_a . By (4), S_a is a clique, so $1 \leq |S_a| \leq 2$ and $S_a \subseteq V(P_u)$. We now break into three cases.

Case 1: $S_b \subseteq V(P_u)$.

By (1) there exist vertices x_a and x_b of P_u such that there exist non adjacent vertices $x_a^* \in S_a^* \cap K_{x_a}$ and $x_b^* \in S_b^* \cap K_{x_b}$.

We first show that there exist such x_a and x_b that are not adjacent. Otherwise, and since $S_a \subseteq V(P_u)$ and $S_b \subseteq V(P_u)$, we have that $S_a \cup S_b = \{x_a, x_b\}$. By replacing x_a and x_b by x_a^* and x_b^* in any principal hole C containing P_u we obtain a path P_C of length $2\ell - 1$, by (3) $S_{\circ}^* \cap P_C = \emptyset$ and so $V(P_C) \cup V(P)$ induces a hole of length at least $2\ell + 2$, a contradiction. So we may assume that x_a and x_b are not adjacent.

Let C be any principal hole of F_0 that contains P_u . By Lemma 9.3, $\{x_a^*, x_b^*\} \cup (V(C) \setminus \{x_a, x_b\})$ induces a hole C^* . Let us apply (5) to C^* , x_a^* and x_b^* . We obtain that the shortest path in C^* between x_a^* and x_b^* has length ℓ or $\ell - 1$. However the path P^* obtained from P_u by replacing x_a by x_a^* and x_b by x_b^* is contained in C^* and it has length at most $\ell - 1$, So wlog, $x_a = u$ and $x_b = u'$. We should then be in the second situation described in (5) and there should exist $z \in S_a^* \cap V(C)$, and $t \in S_b^* \cap V(C)$ such that the shortest path between them on C^* has length $\ell - 1$ and is disjoint from P^* , a contradiction to the assumption that S_a and $S_b \subseteq V(P_u)$.

Case 2: S_b contains a vertex of some principal path P_v distinct from P_u . Up to symmetry, since S_b is a clique (by (4)), we assume that b is anticomplete to $K_{v'}$.

Let y be the vertex of P_u closest to u' such that a has a neighbor $y^* \in K_y$. Let z be the vertex of P_v closest to v such that b has a neighbor $z^* \in K_z$. Possibly y = u' and z = v, but $z \neq v'$ and $y \neq u$ since a has a neighbor in I^* by assumption. In particular, $yz \notin E(G)$. By condition (c) of blowups, $y^*z^* \notin E(G)$.

Let C be the principal hole of F_0 that contains P_u and P_v . By Lemma 9.3, $\{y^*, z^*\} \cup (V(C) \setminus \{y, z\})$ induces a hole C^* . Applying (5) to C^* , y^* and z^* , we obtain that P has length $\ell - 2$ or $\ell - 1$. We denote by P_u^* the path obtained from P_u by replacing y by y^* , and by P_v^* the path obtained from P_v by replacing z by z^* . Let P^* be the path $vP_v^*z^*bPay^*P_u^*u'$ (in case z = v one should replace $vP_v^*z^*$ by z^* , and in case y = u' one should replace $y^*P_u^*u'$ by y^*). The length of P^* is at least ℓ .

Consider now a vertex $r \in A \setminus \{u, v\}$. Depending on the adjacencies of r with v and of r' with u', one of $rvP^*u'w'r'P_rr$ or $rwvP^*u'w'r'P_rr$ or $rwvP^*u'r'P_rr$ or $rvP^*u'r'P_rr$ (with possibly v replaced by z^* in case z = vand u' replaced by y^* in case y = u) is a cycle C_r with at most one possible chord br or bw in case z = v. Hence the length of this cycle should be at most $2\ell + 1$. Since the length of P_r is at least $\ell - 2$ the case where $C_r = rwvP^*u'w'r'P_rr$ cannot occur and we get that at least one of vr, u'r'is an edge of G and so at least one of u, v, r is in A_K . We also notice that if the length of P^* is at least $\ell + 1$ then vr and u'r' should be edges of G, $r \in A_S, u, v \in A_K, z = v, y = u'$ and br is a chord of the cycle $rz^*P^*y^*r'P_rr$. This should be valid for any $r \in A \setminus \{u, v\}$. However since the partition is proper, A_S contains at least 2 elements and we get a contradiction to the fact that S_b is a clique. Hence P^* has length $\ell, y = u', z = v$ and P has length $\ell - 2$.

So by (5), P and C^* form a theta. However, since we have assumed $S_a \cap I \neq \emptyset$, a has a neighbor $t^* \in K_{u^-}$ where u^- is the neighbor of u' in P_u . By Lemma 9.3, since S_a^* is a clique, we have that $\{t^*\} \cup (V(C^*) \setminus \{u^-\})$ induces a hole. This hole and P form a pyramid, a contradiction to (5).

Case 3: We are neither in Case 1 nor in Case 2.

Since we are not in Case 1, S_b contains a vertex of $F_0 \setminus P_u$, and since we are not in Case 2, this vertex must be in $B \cup B'$. Up to symmetry, we assume that $S_b \cap B \neq \emptyset$. Since S_b is a clique (by (4)), $S_b \cap (B' \cup A' \cup I) = \emptyset$. Hence, $S_b \subseteq B \cup \{u\}$ and there exist $x \in B \cap S_b$ and $x^* \in K_x \cap S_b^*$.

Let u_a be the vertex of S_a which is the closest to u in P_u and let u'_a be the vertex of S_a which is the closest to u' in P_u . Notice that, since S_a is a clique (by (3)), either $u_a = u'_a$ or $u_a u'_a$ is an edge. So it may be that $u_a = u$ or $u'_a = u'$ but since $S_a \cap I \neq \emptyset$ we know that $u_a \neq u'$ and $u'_a \neq u$. Let now $u^*_a \in K_{u_a} \cap S^*_a$ and $u'^*_a \in K_{u'_a} \cap S^*_a$. We denote by P^*_u the path obtained from P_u by replacing u_a by u^*_a and, in case $u_a \neq u'_a$, by replacing u'_a by u'^*_a . Notice that if $u_a^* \neq u_a'^*$ then $u_a^* u_a'^* \in E(G)$ since S_a^* is a clique.

Suppose that $u'_a = u^+$, where u^+ is the neighbor of u in P_u . Since H_x contains at least two vertices there exists $v \in H_x \setminus \{u\}$. By (3), by the fact that P contains at least one edge, and because P_u and P_v belong to a hole of F_0 of length 2ℓ in which u^+ and v are at distance at most 3, one of $aPbx^*vP_vv'u'P_u^*u'_a^*a$ or $aPbx^*vP_vv'w'u'P_u^*u'_a^*a$ is a hole of length at least $2\ell + 1$, a contradiction. Hence from now on, we may assume that $u_a \neq u$ (hence a is not adjacent to u) and that if $u_a = u^+$ then $u'_a \neq u_a$. Now by (3) we get that $S_o = \emptyset$.

Suppose first that $u \in H_x$. Let $v \in H_x$ be non-adjacent to u (this is possible since $G[H_x]$ is anticonnected and contains more than one vertex). So, P_u , P_v , x and possibly w' form a hole C (possibly not principal). Let zbe the vertex in $I \cap S_a$ which is the closest to u in P_u and let $z^* \in K_z \cap S_a^*$. By Lemma 9.3, $\{x^*, z^*\} \cup V(C) \setminus \{x, z\}$ induces a hole C^* of $F_1(ux^*, vx^* \in E(G)$ by condition (f) of blowups). The distance between x^* and z^* in C^* is at most $\ell - 1$ since P_u has length at most $\ell - 1$, a contradiction to (5) applied to C^* , x^* and z^* (we cannot have a theta because x^* and z^* are too close in C^* and we cannot have a prism because else b would have a neighbor in K_v , a contradiction to $S_b \subseteq B \cup \{u\}$). Hence, from here on, we may assume that no vertex $x \in B \cap S_b$ is such that $u \in H_x$.

Suppose now that $u \in N(H_x)$. Since $S_b \subseteq B \cup \{u\}$, we have $S_b \cap H_x = \emptyset$. Depending on whether *b* is adjacent to *u* or not, one of $u_a^* a P b u P_u^* u_a^*$ or $u_a^* a P b x^* u P_u^* u_a^*$ (remind that $u_a \neq u$) is a hole, implying that *P* has length at least $\ell - 1$. Let $v \in H_x$, then uv is an edge (since $u \in N(H_x)$). So, $x^* b P a u_a'^* P_u^* u' w' v' P_v v x^*$ or $x^* b P a u_a'^* P_u^* u' v' P_v v x^*$ (in case $u_a'^* = u'$ one should replace $u_a'^* P_u^* u'$ by $u_a'^*$) is a hole of length at least $2\ell + 1$, a contradiction.

Hence, from here on, we may assume that no vertex in $B \cap S_b$ is adjacent to u and so $w \notin B \cap S_b$, in particular $x \neq w$ and x^* is not adjacent to u. Then to avoid a C_4 induced by $\{b, x^*, w, u\}$, b is not adjacent to u and $u_a^* a P b x^* w u P_u^* u_a^*$ is a hole, implying that P has length at least $\ell - 2$. So, for any $v \in H_x$, the hole $x^* b P a u_a'^* P_u^* u' v' P_v v x^*$ or the hole $x^* b P a u_a'^* P_u^* u' w' v' P_v v x^*$ or the hole $x^* b P a u_a'^* P_u^* u' w' v' P_v v x^*$ (in case $u_a' = u'$ one should replace $u_a'^* P^* u'$ by $u_a'^*$) has length 2ℓ if and only if P has length $\ell - 2$, $u_a' = u'$, u' is adjacent to v' and P_v has length $\ell - 2$. This implies that $v \in A_S$ and $u \in A_K$. Furthermore, since $S_a \cap I \neq \emptyset$ and S_a is a clique we get that u_a is the neighbor of u' on the path P_u .

Assume there exists $r \in A_S$ which is adjacent to u. Then $r \notin H_x$ and $rP_r r'w'u'_a aPbx^*wr$ is a hole of length $2\ell + 2$, a contradiction. Hence no such r exists and by setting $\mathbb{A}_K = (A_K \setminus \{u\}) \cup \{x^*\}, \mathbb{A}_S = A_S \cup \{u\}, \mathbb{B} =$

 $\{w\}, \mathbb{A}'_K = (A'_K \setminus \{u'\}) \cup \{a\}, \mathbb{A}'_S = A'_S \cup \{u^*_a\}, \mathbb{B}' = \{u'^*_a\}$ we obtain an even pretemplate partition contained in G with k + 1 paths (the new paths are $uP_u u^*_a$ of length $\ell - 2$ and $x^* bPa$ of length $\ell - 1$), a contradiction.

This proves (6).

(7) We may assume that $S_a \subseteq A \cup B$ and $S_b \subseteq A' \cup B'$.

Otherwise, by (6) and since S_a and S_b are cliques (by (4)), we may assume that $S_a, S_b \subseteq A \cup B$.

We will show that there exists a path Q^* of length at least $2\ell - 2$ whose union with P induces a hole. This is a contradiction because it implies that P has length at most 0. So, to conclude the proof, it remains to prove the existence of Q^* .

By (1), there exist non-adjacent $x_a^* \in S_a^*$ and $x_b^* \in S_b^*$ and for all such x_a^* and x_b^* , $x_a^* a P b x_b^*$ is a path. Let x_a and x_b be the vertices of F_0 such that $x_a^* \in K_{x_a}$ and $x_b^* \in K_{x_b}$. Note that possibly $x_a x_b$ is an edge, but this happens only if $x_a x_b$ is an optional edge of F_0 (since $x_a^* x_b^*$ is not an edge). We break into three cases.

Case 1: $x_a, x_b \in A$.

Then $x_a x_b \notin E(G)$ (otherwise x_a^* and x_b^* would be adjacent) and at least one of x_a, x_b belongs to A_S . So from the definition of templates, there exists a path Q of length $2\ell - 2$ from x_a to x_b consisting in P_{x_a} and P_{x_b} joined either by an edge $x'_a x'_b$ or by a path $x'_a w' x'_b$. By Lemma 9.3, $\{x_a^*, x_b^*\} \cup$ $(V(Q) \setminus \{x_a, x_b\})$ induces the path Q^* that we are looking for. Note that Q^* and P form a hole by (2) and our assumption that $S_a, S_b \subseteq A \cup B$.

Case 2: $x_a \in A$ and $x_b \in B$.

Whether $x_a x_b$ is an optional edge or a non-edge, an immediate consequence of the definition of a template is that there exists a vertex $z \in H_{x_b}$ that is non-adjacent to x_a . By (4), S_a is a clique so $z \notin S_a$. We may furthermore assume that $z \notin S_b$ since else we are in the same situation as in Case 1. By definition of a template, there exists a path Q_0 of length $2\ell - 2$ between x_a and z consisting in P_{x_a} and P_z joined either by an edge $x'_a z'$ or by a path $x'_a w' z'$. Then by Lemma 9.3, $\{x^*_a, x^*_b\} \cup (V(x_b z Q_0 x_a) \setminus \{x_a, x_b\})$ induces a path Q^* of length $2\ell - 1$. Note that Q^* and P form a hole by (2) and our assumption that $S_a, S_b \subseteq A \cup B, z \notin S_b$ and $z \notin S_a$.

Case 3: $x_a, x_b \in B$.

Then $x_a x_b \notin E(G)$ (otherwise x_a^* and x_b^* would be adjacent). Hence, by Lemma 8.4, $H_{x_a} \cup \{x_a\}$ is anticomplete to $H_{x_b} \cup \{x_b\}$. So, let $u_a \in H_{x_a}$ and $u_b \in H_{x_b}$, we may assume that $u_a, u_b \notin S_a \cup S_b$ since else we are in the situation of Case 1 or 2. By definition of a template, there exists a path Q_0 of length $2\ell - 2$ between u_a and u_b consisting in P_{u_a} and P_{u_b} joined either by an edge $u'_a u'_b$ or by a path $u'_a w' u'_b$. By Lemma 9.3, $Q^* = x^*_a u_a Q_0 u_b x^*_b$ is also a path, it is of length 2ℓ . Now, it is easy to verify that Q^* and P form a hole of length more than 2ℓ , a contradiction.

This proves (7).

(8) $S_{\circ} = \emptyset$.

By (4) and (7), if $S_{\circ} \neq \emptyset$, then $\ell = 4$, and there exists a principal path $P_u = ucu'$ of F_0 such that $S_a = \{u\}$, $S_b = \{u'\}$ and $S_{\circ} = \{c\}$. Let $u^* \in K_u \cap S_a^*$, $c^* \in S_{\circ}^*$ and $u'^* \in K_{u'} \cap S_b^*$. Observe that by (2), each of u^*c^*, u'^*c^* is an edge and by definition, c^* has a neighbor in int(P).

Let $P_v = v \dots v'$ be any principal path distinct from P_u . Now, P_v , P, u^* , u'^* , c^* and possibly w and/or w' form a proper wheel centered at c^* , a contradiction to Lemma 3.10. This proves (8).

 $(9) \ (S_a \cup S_b) \cap (A_S \cup A'_S) = \emptyset.$

Notice that each of $S_a \cap A_S$ and $S_b \cap A'_S$ contains at most one vertex since S_a and S_b are cliques.

Case 1: S_a contains $x \in A_S$ and S_b contains $y' \in A'_S$.

Let $x^* \in K_x \cap S_a^*$ and $y'^* \in K_{y'} \cap S_b^*$. We denote by P_x^* (resp. P_y^*) the path obtained from P_x (resp. P_y) by replacing x (resp. y') by x^* (resp. y'^*). If x = y then $ax^*P_x^*x'bPa$ is a hole, so P has length ℓ . Since the partition is proper there exists an other vertex $r \in A_S$ and $ax^*wrP_rr'w'y'^*bPa$ is a cycle of length $2\ell + 4$ whose possible chords are aw and bw', in any case the cycle contains a too long hole. Hence $x \neq y$.

Assume that $aw, bw' \notin E(G)$. Then $ax^*P_x^*x'w'y'^*bPa$ is a hole and hence P has length $\ell - 2$. We know that the template has at least one more principal path P_r . In case there exists $r \in A_S$ then $ax^*wrP_rr'w'y'^*bPa$ is a hole of length $2\ell + 2$, a contradiction. So any $r \in A \setminus \{x, y\}$ belongs to A_K . If rx and ry are in E(G) then $ax^*rP_rr'w'y'^*bPa$ is a cycle of length $2\ell + 2$ whose only possible chord is ar, so we get a contradiction. So by symmetry we get that r is adjacent to exactly one of x and y and r' is adjacent to exactly one of x' and y'. In case r is adjacent to y and not to xthen $ax^*wrP_rr'w'y'^*bPa$ is a hole of length $2\ell + 3$, a contradiction. So we may assume that r is adjacent to x and not to y (and then r' is adjacent to y' and not to x'), and this is true for any $r \in A_K$.

Let x^+ be the neighbour of x in P_x and y^- be the neighbour of y' in P_y . By setting $\mathbb{A}_K = A_K, \mathbb{A}_S = \{a, x^+, w\}, \mathbb{B} = \{x^*\}, \mathbb{A}'_K = A'_K, \mathbb{A}'_S = \{a, x^+, w\}, \mathbb{B} = \{x^*\}, \mathbb{A}'_K = A'_K, \mathbb{A}'_S = \{x^*\}, \mathbb{A}'_K = \{x^*\}, \mathbb{A}'_K$

 $\{b, w', y^-\}, \mathbb{B}' = \{y'^*\}$ we obtain an even pretemplate partition contained in G with k + 1 paths (the new paths are $aPb, x^+P_xx'w', wyP_yy^-$ each of length $\ell - 1$), a contradiction. Hence aw or bw' is an edge of G.

So by symmetry we may set that $aw \in E(G)$. Then $awyP_y^*y'^*bPa$ is a hole, so P has length $\ell - 1$. If $bw' \notin E(G)$ then $ax^*P_x^*x'w'y'^*bPa$ is a hole of length $2\ell + 1$, a contradiction. So $bw' \in E(G)$. Suppose A_S contains a vertex $r \neq x, y$ then $awrP_rr'w'y'^*bPa$ is hole of length $2\ell + 1$, a contradiction. Hence $A_S = \{x, y\}$ and $ar, br' \in E(G)$ for every $r \in A_K$.

So, by setting $\mathbb{A}_K = A_K \cup \{a\}, \mathbb{A}_S = \{x^*, y\}, \mathbb{B} = \{w\}, \mathbb{A}'_K = A'_K \cup \{b\}, \mathbb{A}'_S = \{x', y'^*\}, \mathbb{B}' = \{w'\}$ we obtain an even pretemplate partition contained in G with k + 1 paths (the new paths are aPb of length $\ell - 1$ and P_x^*, P_y^* of length $\ell - 2$), a contradiction.

Case 2: $S_a \cap A_S = \emptyset$, S_a contains a vertex $x \in A_K$ and S_b contains a vertex $y' \in A'_S$ (symmetric to the case where $S_b \cap A'_S = \emptyset$, S_b contains a vertex $y' \in A'_K$ and S_a contains a vertex $x \in A_S$).

Let $x^* \in K_x \cap S_a^*$ and $y'^* \in K_{y'} \cap S_b^*$. Suppose that $xy \in E(G)$. Then $ax^*yP_y^*y'^*bPa$ is a hole and hence P has length $\ell-1$. Then $ax^*P_x^*x'w'y'^*bPa$ is a cycle with at most one possible chord (bw') of length $2\ell+2$, a contradiction. So $xy \notin E(G)$ and consequently $x'y' \in E(G)$. Since x^* and y'^* both belong to the hole $x^*P_xx'y'^*P_y^*ywx^*$, by (5) we get that P and this hole induce either a theta or a prism.

- Subcase 2.1: $x^*P_x^*x'y'^*P_y^*ywx^*$ and P induce a theta.

Then P has length $\ell - 2$ and $aw, bx' \notin E(G)$. Assume that y has a neighbour v in A. Then since $y \in A_S$ we have that $v \in A_K$ and hence $v'y'^* \notin E(G)$ and $xv, x'v' \in E(G)$. Hence to avoid a C_4 induced by $\{b, x', y'^*, v'\}$ it should be that $bv' \notin E(G)$. So $aPby'^*x'v'P_vvxa$ is a cycle of length $2\ell + 2$ whose only possible chord is va, a contradiction to the fact that G belongs to $C_{2\ell}$. We may then conclude that y has no neighbor in A and so by setting $\mathbb{A}_K = (A_K \setminus \{x\}) \cup \{x^*, w\}, \mathbb{A}_S = (A_S \setminus \{y\}) \cup \{a\}, \mathbb{B} = \emptyset, \mathbb{A}'_K =$ $A'_K \cup \{y'^*\}, \mathbb{A}'_S = (A'_S \setminus \{y'\}) \cup \{b\}, \mathbb{B}' = B'$, we obtain an even pretemplate partition contained in G with k + 1 paths (the new paths are aPb of length $\ell - 2$ and $x^*P_x^*x'$ and $wyP_y^*y'^*$ of length $\ell - 1$, \mathbb{A}_S and \mathbb{A}'_S are both stable sets), a contradiction.

- Subcase 2.2: $x^*P_x^*x'y'^*P_y^*ywx^*$ and P induce a prism. Then P has length $\ell - 1$ and $aw, bx' \in E(G)$.

So, by setting $\mathbb{A}_K = (A_K \setminus \{x\}) \cup \{x^*, a\}, \mathbb{A}_S = A_S, \mathbb{B} = B, \mathbb{A}'_K = A'_K \cup \{b\}, \mathbb{A}'_S = (A'_S \setminus \{y'\}) \cup \{y'^*\}, \mathbb{B}' = B'$ we obtain an even pretemplate

partition contained in G with k + 1 paths (the new paths are aPb, $x^*P_x^*x'$ and $yP_y^*y'^*$ of lengths respectively $\ell - 1$, $\ell - 1$, $\ell - 2$), a contradiction.

Case 3: $S_a \cap A = \emptyset$, S_a contains a vertex $x \in B$ and S_b contains a vertex $y' \in A'_S$ (symmetric to the case where $S_b \cap A' = \emptyset$, S_b contains a vertex $y' \in B'$ and S_a contains a vertex $x \in A_S$).

If $xy \in E(G)$ then because of the hole $ax^*yP_yy'^*bPa$ the path P has length $\ell - 1$ and once again it is enough to replace x by x^* , y' by y'^* and add a to A_K and b to A'_K in order to obtain an ℓ -pretemplate partition with k + 1 principal paths, hence $xy \notin E(G)$. So, we may assume that a is not adjacent to w. Then because of $ax^*wyP_y^*y'^*bPa$, the path P has length $\ell - 2$.

So, by setting $\mathbb{A}_K = A_K \cup \{x^*\}, \mathbb{A}_S = A_S, \mathbb{B} = \{w\}, \mathbb{A}'_K = A'_K \cup \{b\}, \mathbb{A}'_S = (A'_S \setminus \{y'\}) \cup \{y'^*\}, \mathbb{B}' = B'$ we obtain an even pretemplate partition contained in G with k + 1 paths (the new paths are x^*aPb of length $\ell - 1$ and $yP_y^*y'^*$ of length $\ell - 2$), a contradiction.

This proves (9).

We may now conclude the proof. By (9) we have that $S_a \subseteq A_K \cup B$ and $S_b \subseteq A'_K \cup B'$. Notice that if the length of P is $\ell - 1$ (resp. $\ell - 2$) then we get a contradiction since we can add a to A_K and b to A'_K (resp. a to A_S and b to A'_S) and replace any vertex $x \in S_a$ and any vertex $y' \in S_b$ by respectively $x^* \in K_x \cap S^*_a$ and $y'^* \in K_y \cap S^*_b$ in order to get an ℓ -pretemplate partition with one more principal path. So from now on we may assume that the length of P is neither $\ell - 1$ nor $\ell - 2$.

Let us assume that there exist $x \in S_a \cap A_K$ and $y' \in S_b \cap A'_K$ and let $x^* \in K_x \cap S_a^*$ and $y'^* \in K'_y \cap S_b^*$. If x = y then $ax^*P_x^*y'^*bPa$ is a hole, and hence P has length $\ell-1$, a contradiction. So we may now assume that $x \neq y$, $S_a \cap K_y = \emptyset$ and $S_b \cap K_{x'} = \emptyset$. Then because of the hole $ax^*yP_y^*y'^*bPa$ the length of P is $\ell-2$, a contradiction again.

So from now on we may assume by symmetry that $S_a \cap A_K = \emptyset$. Then there exists $x^* \in K_x \cap S_a^*$ for some $x \in B$. Assume that there exists $y'^* \in K'_y \cap S_b^*$ for some $y' \in A_K$. In case $xy \in E(G)$ then P has length $\ell-2$ because of the hole $ax^*yP_y^*y'^*bPa$ and we would get a contradiction. Hence $xy, aw \notin E(G)$. Now the hole $ax^*wyP_y^*y'^*bPa$ implies that P has length $\ell-3$. Then any vertex $u \in A$ which is adjacent to x, is adjacent to y: if $u \in A_K$ by definition of an even template, if $u \in A_S$ because else $ax^*uP_uu'y'^*bPa$ is hole of length $2\ell - 1$. However, as any vertex in B, x has two non adjacent neighbors $r, s \in A$ and so G contains a square x^*rysx^* , a contradiction. So from now on we may assume that $S_b \subseteq B'$ and there exists $y'^* \in K_{y'} \cap S_b^*$ for some $y' \in B'$. By definition, x has at least one neighbor r in A_S . If $y'r' \in E(G)$ then because of the hole $ax^*rP_rr'y'^*bPa$ we get that P has length $\ell - 2$, a contradiction. So $r'y' \notin E(G)$ and then $y' \neq w'$, $bw' \notin E(G)$ and P should have length $\ell - 3$, because of the cycle $ax^*rP_rr'w'y'^*bPa$. By symmetry, we also have $x \neq w$.

So, by setting $\mathbb{A}_K = A_K \cup \{x^*\}, \mathbb{A}_S = A_S, \mathbb{B} = \{w\}, \mathbb{A}'_K = A'_K \cup \{y^*\}, \mathbb{A}'_S = A'_S, \mathbb{B}' = \{w'\}$ we obtain an even pretemplate partition contained in G with k + 1 paths (the new path is x^*aPby^* of length $\ell - 1$), a contradiction.

9.6 End of the proof

We may now conclude the proof of Lemma 9.1. If $G \setminus F_1$ is empty, then conclusion (a) holds. If $G \setminus F_1$ is non-empty and $G \setminus F_2$ is empty, then conclusion (b) holds. Otherwise, we consider a connected component D of $G \setminus F_2$ and apply Lemma 9.15. We then see that G has a clique cutset, so conclusion (c) holds.

10 Proof of Theorem 10.1

Theorem 10.1 Let $\ell \geq 4$ be an integer. If G is a graph in $C_{2\ell}$ then one of the following holds:

- (a) G is a ring of length 2ℓ ;
- (b) G is a proper blowup of a twinless even ℓ -template;
- (c) G has a universal vertex or
- (d) G has a clique cutset.

Proof. By Lemma 3.9, G contains no unbalanced prism, no unbalanced theta, no pyramid and no proper wheel. Also, clearly G contains no C_4 and no C_5 . Hence, by Theorem 3.13, we may assume that G contains a prism or a theta for otherwise one of the conclusions (a), (c) or (d) holds. The result then follows from Lemma 9.1.

References

- Eli Berger, Paul Seymour, and Sophie Spirkl. Finding an induced path that is not a shortest path. *Discrete Mathematics*, 344(7):112398.1– 112398.6, 2021.
- [2] Valerio Boncompagni, Irena Penev, and Kristina Vušković. Cliquecutsets beyond chordal graphs. *Journal of Graph Theory*, 91(2):192–246, 2019.
- [3] Yung-Chung Chiu and Hsueh-I Lu. Blazing a Trail via Matrix Multiplications: A Faster Algorithm for Non-shortest Induced Paths. arXiv:2109.15268, 2022.
- [4] M. Chudnovsky, N. Robertson, P. Seymour, and R. Thomas. The strong perfect graph theorem. Annals of Mathematics, 164(1):51–229, 2006.
- [5] Václav Chvátal and Peter L. Hammer. Aggregation of inequalities in integer programming. In P.L. Hammer, E.L. Johnson, B.H. Korte and G.L. Nemhauser, editors, *Computational Graph Theory, Computing Supplementum, Annals of Discrete Mathematics*, 1, pages 145–162. North-Holland, 1975.
- [6] M. Conforti, G. Cornuéjols, A. Kapoor, and Kristina Vušković. Universally signable graphs. *Combinatorica*, 17(1):67–77, 1997.
- [7] Linda Cook. On recognition algorithms and structure of graphs with restricted induced cycles. PhD thesis, Princeton University, May 2021.
- [8] Linda Cook, Jake Horsfield, Myriam Preissmann, Cléophée Robin, Paul Seymour, Ni Luh Dewi Sintiari, Nicolas Trotignon, and Kristina Vušković. Graphs with all holes the same length. arXiv:2110.09970, 2022.
- M.V.G. da Silva and Kristina Vušković. Decomposition of even-hole-free graphs with star cutsets and 2-joins. *Journal of Combinatorial Theory*, Series B, 103(1):144–183, 2013.
- [10] G.A. Dirac. On rigid circuit graphs. Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg, 25:71–76, 1961.
- [11] Stéphane Foldes and Peter L. Hammer. Split graphs. In Proceedings of the Eighth Southeastern Conference on Combinatorics, Graph Theory

and Computing (Louisiana State Univ., Baton Rouge, La.), pages 311–315, 1977. Congressus Numerantium, No. XIX, Utilitas Math., Winnipeg, Man.

- [12] Angèle M. Foley, Dallas J. Fraser, Chính T. Hoàng, Kevin Holmes, and Tom P. LaMantia. The intersection of two vertex coloring problems. *Graphs Comb.*, 36(1):125–138, 2020.
- [13] Martin Charles Golumbic. Trivially perfect graphs. Discrete Mathematics, 24(1):105–107, 1978.
- [14] Chính Hoàng and Nicolas Trotignon. A class of graphs of large rankwidth. CoRR, abs/2007.11513, 2020.
- [15] Frédéric Maffray, Irena Penev, and Kristina Vušković. Coloring rings. Journal of Graph Theory, 96(4):642–683, 2021.
- [16] Irena Penev. On the clique-width of $(4K_1, C_4, C_5, C_7)$ -free graphs. Discrete Applied Mathematics, 285:688–690, 2020.
- [17] Cléophée Robin. *Hereditary classes of graphs: from structure to coloring.* PhD thesis, Université Grenoble Alpes, October 2021.
- [18] D. Seinsche. On a property of the class of *n*-colorable graphs. Journal of Combinatorial Theory, Series B, 16(2):191–193, 1974.
- [19] Kristina Vušković. The world of hereditary graph classes viewed through Truemper configurations. In S. Gerke S.R. Blackburn and M. Wildon, editors, *Surveys in Combinatorics, London Mathematical Society Lecture Note Series*, volume 409, pages 265–325. Cambridge University Press, 2013.
- [20] E. S. Wolk. A note on the comparability graph of a tree. Proceedings of the American Mathematical Society (1 ed.), 16:17–20, 1965.